

The Coronary Angiography and prevalence of Renal Artery Stenosis at Tertiary Care Center.

Avanish kumar Bhardwaj¹, Jayesh Sharma¹

¹DM Final Year Cardiology, Department of Cardiology, S.P. Medical College, Bikaner.

Received: February 2020

Accepted: February 2020

ABSTRACT

Background: Renal Artery Stenosis (RAS) is a self-determining risk factor for cardiovascular diseases. The present study was designed to assess the prevalence of renal artery stenosis in patients with Coronary Artery Disease (CAD) who underwent Coronary Angiography (CAG). **Methods:** The consecutive CAD patients undergoing CAG and renal angiography were studied from December 2018 to December 2019. The presence of risk factors such as age, hypertension, diabetes, left ventricular function and myocardial infarction were assessed. The degree of arterial stenosis was categorised into mild, moderate and severe and at least 50% narrowing of the arterial lumen was considered as arterial stenosis. Data was analysed by using SPSS 20.0 software. **Results:** Out of 878 patients, the prevalence of RAS was calculated as 33 (3.8%) patients, the majority of them were male (75.9%). Out of 33 patients, 19 (57.6%), 6 (18.2%) and 8 (24.2%) patients were classified as mild, moderate and severe RAS, respectively. Using multiple variables including age, sex, hypertension, diabetes mellitus, multivessel disease and left ventricular dysfunction patients were considered as predictors of RAS. The significant number of a patient had unilateral 18 (54.5%) and 11 (33.3%) bilateral RAS observed from renal angiography. **Conclusion:** The present study suggests that the renal angiography in combination with coronary artery angiography in CAD patients provides the opportunity for identification of RAS.

Keywords: Coronary Artery Disease; Renal Angiography; Renal Artery Stenosis.

INTRODUCTION

Renal Artery Stenosis (RAS) was distinct as any vascular lesion causing narrowing of the renal artery thereby impairing blood flow to the kidney. RAS is frequently present without any clinical signs or symptoms and is one of the etiological factors for renal insufficiency and hypertension and atherosclerosis.^[1] Consequence of Atherosclerotic Renal Artery Stenosis (ARAS) disease leads to renal atrophy over a period of time and chronic kidney disease despite the control of hypertension. Angioplasty of ARAS with renal artery stenting is a beneficial treatment strategy to restore and preserve renal function and to control blood pressure.^[2]

There is a linear relationship between RAS and severity of CAD, but act as an independent predictor of mortality in CAD patients. The treatment of RAS would act as a remedy for severe hypertension and ischaemic nephropathy. The mortality rate in patients with cardiovascular disease increases in the presence of RAS. The occurrence of RAS ranges from 3-30% in CAD patients with varying severity that underwent Coronary Angiography (CAG). However, RAS must be carefully observed, because most patients with RAS have no signs or symptoms.^[3,4] The presence of ARAS exacerbates the CAD and course

of the medical condition. There was a strong relation between systemic hypertension and RAS because of pathophysiologic stimulus, including activation of the renin-angiotensin system by RAS, hypertension triggered by ischaemic nephropathy and aggravation of renal atherosclerosis by hypertension.^[5,6]

Hence, performing the renal angiography reduced the progression of RAS during Coronary Angiography (CAG) as it can be safe and cost-effective diagnostic approach in patients with CAD.^[3] However, regular assessment of RAS in asymptomatic patients suffering from CAD is difficult to endorse, because of the lack of experience of clinical benefits related to renal artery intervention in patients with RAS.

MATERIALS AND METHODS

From December 2018 to december 2019 renal angiography was performed in patients undergoing coronary angiography at Department of Cardiology, S.P.Medical College Bikaner India.

Patient Selection-

This study included male or non-pregnant female patients ≥ 18 years of age with a diagnosis of angina equivalents of class III or IV severity, angina of less severity, but having positive treadmill exercise stress test at low workloads or delayed recovery of depressed ST segments, atypical chest pain syndromes. Additional eligibility criteria were the presence of post myocardial infarction patients who are positive for inducible ischaemia on exercise stress test.

Name & Address of Corresponding Author

Dr. Jayesh Sharma,
DM Final Year Cardiology,
Department of Cardiology,
S.P. Medical College,
Bikaner.

Procedure-

All patients were subjected to a pre-procedure clinical evaluation, which included a thorough history and physical examination pertaining to the cardiovascular system. ECGs were taken for all eligible patients and were subjected to a treadmill exercise stress test on the standard Bruce or modified Bruce protocol. Echocardiographic evaluation was done for all to assess regional wall motion abnormality and Left Ventricular Function (LV). Informed consent from all participants was obtained before the procedure. Coronary angiography was performed preferably with Judkins left and right coronary catheters of appropriate size. After completion of coronary angiography, the Judkins right catheter was withdrawn into the abdominal aorta and renal arteries were selectively cannulated. Diameter stenosis in percentage was recorded. Less than 50% stenosis was classified as mild; 50-75% as moderate and >75% as severe. The incidence of renal artery stenosis and its relation to gender, prevalence of hypertension and CAD were determined. In addition, the incidence of RAS in positive treadmill exercise stress test population was estimated.

RESULTS

Table 1: Baseline and Frequency Distribution of Prevalence of Renal Artery Stenosis among Patients Undergoing Coronary Angiography

Renal Stenosis (n=33)			
Variables	Mild (<50%)	Moderate (50-75%)	Severe (>75%)
MALE	12 (36.3%)	5 (15.2%)	6 (18.2%)
FEMALE	7 (21.2%)	1 (3.0%)	2 (6.1%)
Age			
<50	6 (18.2%)	1 (3.0%)	3 (9.1%)
51-60	6 (18.2%)	2 (6.1%)	1 (3.0%)
>60	7 (21.2%)	3 (9.1%)	4 (12.1%)
Disease Condition			
Diabetic	7 (21.2%)	1 (3.0%)	1 (3.0%)
Non-diabetic	12 (36.3%)	5 (15.2%)	7 (21.2%)
Hypertensive	5 (15.2%)	3 (9.1%)	5 (15.2%)
Non-hypertensive	14 (42.4%)	3 (9.1%)	3 (9.1%)
CAD			
LMCA	1 (3.0%)	1 (3.0%)	0 (0%)
SVD	1 (3.0%)	0 (0%)	0 (0%)
DVD	9 (27.2%)	1 (3.0%)	3 (9.1%)
TVD	7 (21.2%)	5 (15.2%)	2 (6.1%)
MILD	2 (6.1%)	0 (0%)	1 (3.0%)
NORMAL	0 (0%)	1 (3.0%)	2 (6.1%)
LV Function			
Normal	15 (45.5%)	5 (15.2%)	4 (12.1%)
Dysfunction	4 (12.1%)	1 (3.0%)	3 (9.1%)
Myocardial Infarction			
IWMI	4 (12.1%)	2 (6.1%)	2 (6.1%)
AWMI	7 (21.2%)	1 (3.0%)	2 (6.1%)
Type of Stenosis			
Unilateral	15 (45.5%)	3 (9.1%)	0 (0%)
Bilateral	3 (9.1%)	4 (12.1%)	4 (12.1%)
Branch	1 (3.0%)	1 (3.0%)	2 (6.1%)
Site of Involvement			
Left renal artery	5 (15.2%)	3 (9.1%)	2 (6.1%)
Right renal artery	11 (33.3%)	3 (9.1%)	3 (9.1%)
Bilateral	3 (9.1%)	0 (0%)	3 (9.1%)




Figure 1: Schematic Representation of Disposition of Subjects who Underwent Renal Angiogram to Detect Renal Artery Stenosis.




Figure 2: Association of Positive Stress Test in Renal Artery Stenosis Patients




Figure 3: Angiogram of Left Renal Artery Showing Ostial Stenosis (Arrows)

A total of 1101 patients with CAD proven by CAG and consecutive CAG conducted over a period from December 2018 to december2019 were studied. Renal angiograms were done in 878 patients. Most of the patients were males (666; 75.9%). The disposition

of patients who underwent renal angiogram to detect RAS is outlined in [Figure 1].

Renal artery stenosis was detected in 33 (3.8%) patients [Table 1]. There were 10 (30.3%) women and 23 (69.7%) men. The RAS was further graded as mild (<50%), moderate (50-70%) and severe (>75%). Among 33 patients, 19 (57.6%), 6 (18.2%), 8 (24.2%) subjects have having mild, moderate and severe RAS, respectively. Out of 33 patients, 9 (27.3%) cases were diabetic and 24 (72.7%) were non-diabetic patients. Whereas, 13 (39.4%) were hypertensive and 20 (60.6%) were non-hypertensive patients. Among 13 hypertensive patients, 9.1% had moderate RAS and 15.2% had severe RAS. Among 33 patients, only 3 had normal coronary arteries. The prevalence of RAS for Single Vessel Disease (SVD), Left Main Coronary Artery (LMCA), Double Vessel Disease (DVD) and Triple-Vessel Disease (TVD) were in 1, 2, 13 and 14, respectively. Approximately, one-quarter of 9 (27.3%) patients had an incidence 57.6 LVdysfunction and 19 (57.6%) had a prior myocardial infarction. Of 33 patients, 18 (54.5%) individual had unilateral RAS followed by 11 (33.3%) patients suffering from bilateral RAS. Furthermore, positive treadmill exercise stress test (20; 60.6%) was represented notably higher in patients suffering from RAS [Figure 2]. Out of 33 RAS participants, 30 (91%) cases illustrated the lesion was ostial [Figure 3]. Amongst those with mild stenosis, 5 (15.2%) had involvement of left renal artery, 11 (33.3%) had involvement of right renal artery and 3 (9.1%) had bilateral involvement. Moderate stenosis was observed only in 3 patients on either side of the kidney.

DISCUSSION

Coronary artery disease is hardening and narrowing of arteries that supply blood to heart muscle. It is commonly associated with atherosclerosis of cerebrovascular, peripheral and renal arteries. Some of the patients suffering from CAD may also have presence of RAS. Hence, there is need of a cardiologist to undertake a more global approach to patients with CAD.

The present study was conducted to determine the prevalence of RAS in patient population referred for diagnostic CAG suffering from CAD and its multiple variables were based on number of diseased coronary arteries, age, sex, history of systemic hypertension and diabetes mellitus, LV functions, type of myocardial infarction, type of stenosis and site of involvement. In this study, we have investigated 878 CAD patients who underwent renal angiography and found that the prevalence of RAS in this population was 33 (3.76%), which was relatively low as reported elsewhere.^[6,7] The reported prevalence was usually between 14% and 20% depending on the study inclusion criteria, sample size and severity of renal stenosis criteria, presence or absence of peripheral

arterial disease, serum creatinine concentration and presence of triple-vessel CAD or history of coronary artery bypass graft surgery.^[8,9] The present findings suggest that males were more commonly affected by ARAS.

A majority of the study population had unilateral 18 (54.5%) RAS, while 11 (33.3%) of them suffered from bilateral RAS and only 4 (12.1%) had branch RAS. On the other hand, previously published prospective study revealed that 30% of patients diagnosed with RAS underwent coronary angiography of which 15% demonstrated that significant stenosis (11% unilateral and 4% bilateral).^[10] The unilateral RAS was associated with increased renin and aldosterone levels that leads to vasoconstriction and increased peripheral vascular resistance. Furthermore, in bilateral RAS, both the kidneys activated renin and aldosterone flow causing retention of water and sodium to restore the volume and maintain hypertension.^[11]

Diabetes and hypertension, common risk factors for atherosclerosis were not mainly related with significant RAS, because they were already reflected by other variables, specifically significant old age, CAD and LV dysfunction. The high predominance of hypertension in patients experiencing CAG also presented significant association between RAS and hypertension (39.4%). While in previously published literature, Rimoldi et al reported lower percentage of RAS (8%) in hypertensive patients.^[12] On the contrary, prevalence of ARAS was estimated 13% as reported by Yamashita et al.^[13] Accordingly, this discrepancy between published literatures partially correlated with the ethnic, regional and lifestyle pattern differences.

The prevalence of RAS in males was more commonly affected by atherosclerotic RAS and it was found to be significantly high in the patients with diabetes mellitus. Our study confirmed that age >60 years, male sex, multivessel disease and LV dysfunction patients were predictors of RAS.^[14] The vast majority of patients with RAS had a positive exercise stress tests and incidence of LV dysfunction was more with increase in severity of RAS. These results are in agreement with study by Tumelero et al in which they concluded the presence of a strong relationship between LV dysfunction and RAS.^[15]

For clinical investigation of CAD patients, it is difficult to perform renal angiography in all selected patients because of the lack of evidence of benefit and low prevalence of RAS. Hence, a proper differentiation should be made in patient with considerable RAS who are at higher risk of cardiovascular events and need close surveillance. As in CAD patients, RAS is an independent predictor of mortality.^[16] So, from the observations of cardiologist, it advisable to perform renal angiography at the time of CAG in selected patients undergoing CAG.

CONCLUSION

The present finding demonstrated that prevalence of renal artery stenosis observed in patients with coronary artery disease who underwent coronary angiography.

REFERENCES

1. Payami B, Jafarizade M, Beladi Mousavi SS, et al. Prevalence and predictors of atherosclerotic renal artery stenosis in hypertensive patients undergoing simultaneous coronary and renal artery angiography; a cross-sectional study. *J Renal Inj Prev* 2016;5(1):34-38.
2. Gonçalves JAA, Amorim JE, Neto MMS, et al. Eficácia clínica da revascularização renal percutânea com implante de stent em pacientes com doença renovascular aterosclerótica. *Arq Bras Cardiol* 2007;88(1):85-90.
3. White CJ, Jaff MR, Haskal ZJ, et al. Indications for renal arteriography at the time of coronary arteriography: a science advisory from the American Heart Association Committee on diagnostic and interventional cardiac catheterization, council on clinical cardiology, and the councils on cardiovascular radiology and intervention and on kidney in cardiovascular disease. *Circulation* 2006;114(17):1892-1895.
4. Garovic VD, Textor SC. Renovascular hypertension and ischemic nephropathy. *Circulation* 2005;112(9):1362-1374.
5. Schiffrin EL, Lipman ML, Mann JF. Chronic kidney disease: effects on the cardiovascular system. *Circulation* 2007;116(1):85-97.
6. Ollivier R, Boulmier D, Veillard D, et al. Frequency and predictors of renal artery stenosis in patients with coronary artery disease. *Cardiovasc Revasc Med* 2009;10(1):23-29.
7. Lee Y, Shin JH, Park HC, et al. A prediction model for renal artery stenosis using carotid ultrasonography measurements in patients undergoing coronary angiography. *BMC Nephrol* 2014;15:60.
8. Bageacu S, Cerisier A, Isaaz K, et al. Incidental visceral and renal artery stenosis in patients undergoing coronary angiography. *Eur J Vasc Endovasc Surg* 2011;41(3):385-390.
9. Ozkan U, Oguzkurt L, Tercan F, et al. The prevalence and clinical predictors of incidental atherosclerotic renal artery stenosis. *Eur J Radiol* 2009;69(3):550-554.
10. Harding MB, Smith LR, Himmelstein SI, et al. Renal artery stenosis: prevalence and associated risk factors in patients undergoing routine cardiac catheterization. *J Am Soc Nephrol* 1992;2(11):1608-1616.
11. Liard JF, Cowley AW, McCaa RE, et al. Renin, aldosterone, body fluid volumes, and the baroreceptor reflex in the development and reversal of Goldblatt hypertension in conscious dogs. *Circ Res* 1974;34(4):549-560.
12. Rimoldi SF, de Marchi SF, Windecker S, et al. Screening renal artery angiography in hypertensive patients undergoing coronary angiography and 6-month follow-up after ad hoc percutaneous revascularization. *J Hypertens* 2010;28(4):842-847.
13. Yamashita T, Ito F, Iwakiri N, et al. Prevalence and predictors of renal artery stenosis in patients undergoing cardiac catheterization. *Hypertens Res* 2002;25(4):553-557.
14. Cohen MG, Pascua JA, Garcia-Ben M, et al. A simple prediction rule for significant renal artery stenosis in patients undergoing cardiac catheterization. *Am Heart J* 2005;150(6):1204-1211.
15. Tumelero RT, Duda NT, Tognon AP, et al. Prevalence of renal artery stenosis in 1,656 patients who have undergone cardiac catheterization. *Arq Bras Cardiol* 2006;87(3):248-253.
16. Mui KW, Sleeswijk M, van den Hout H, et al. Incidental renal artery stenosis is an independent predictor of mortality in

patients with peripheral vascular disease. *J Am Soc Nephrol* 2006;17(7):2069-2074.

Copyright: © Annals of International Medical and Dental Research. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bhardwaj A, Sharma J. The Coronary Angiography and Pprevalence of Renal Artery Stenosis at Tertiary Care Center. *Ann. Int. Med. Den. Res.* 2020; 6(3):ME39-ME42.

Source of Support: Nil, **Conflict of Interest:** None declared