

Prevalence and Management of Postpartum Anaemia in a Tertiary Care Hospital in North India

Preeti Sharma¹

¹Assistant Professor, Department of Obstetrics & Gynaecology, Muzaffarnagar Medical College, Uttar Pradesh, India.

Received: March 2019

Accepted: March 2019

Copyright: © the author(s), publisher. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Postpartum iron deficiency anaemia (IDA) is common in women in resource-poor countries. Most women are treated with either oral iron or blood transfusion. The aim of our study was to find out the prevalence of postpartum anaemia; and to compare the effect of treatment with either oral ferrous sulphate or intravenous iron sucrose on postpartum IDA. **Methods:** 102 postpartum women with proven iron deficiency anaemia with hemoglobin <9gm/dl and serum ferritin <15 µgm/l were included in the study. They were randomized to receive either oral ferrous sulphate 200 mg twice daily for 6 weeks (group 1) or intravenous ferrous sucrose 200 mg, two to three doses given on alternate days (group 2). Total iron deficit was calculated using standard formula and results analyzed. Main outcome measures: Hemoglobin, hematocrit, red cell indices and ferritin were measured on day 2-3, 1-2 weeks and 6 weeks postpartum. **Result:** The prevalence of postpartum anaemia was 19.92%. By 1-2 weeks, hemoglobin level in women treated with intravenous iron had risen from 7.81 ± 0.849 to 9.88 ± 0.760 gm/dl which was more than those treated with oral iron ($p < 0.0001$); although by 6 weeks, there was no significant difference between the two groups. Ferritin levels rose rapidly in those treated with intravenous iron and remained significantly higher than in those treated with oral iron ($p < 0.0001$). **Conclusion:** Intravenous iron sucrose increases the hemoglobin level more rapidly than oral ferrous sulphate in women with postpartum IDA. It also replenishes iron stores more rapidly.

Keywords: Postpartum Iron deficiency anaemia, Ferritin, Iron sucrose, ferrous sulphate.

INTRODUCTION

Postpartum anaemia is a common problem throughout the world and for most women is self-limiting, resolving within a week. For some women however, particularly in resource-poor countries, it is a major cause of maternal morbidity and mortality.^[1] Postpartum haemoglobin (Hb) levels of <10 g/dl are observed in up to 30% of women, with more severe anaemia (Hb < 8g/dl) seen in 10%. Iron deficiency is the principal cause. This is partly attributable to an iron deficit during pregnancy, caused by the increased iron demands of the fetoplacental unit and an increased maternal red cell mass. Irrespective of mode of delivery, blood loss is a contributing factor, with 5% of deliveries involving loss of more than 1L. It appears to be higher in unfavorable socioeconomic conditions.^[2]

The symptoms of postpartum anaemia vary and may include breathlessness, fatigue, palpitations, dizziness, maternal infections particularly of the urinary tract, lactation failure and prolonged hospital

stay depending on the severity of blood loss and related anaemia. It has also been shown to be strongly associated with depression, stress and cognitive function in the postpartum period and may result in difficulty for the mother to care for her baby, thereby influencing the emotional mother-infant bond.

The standard approach to treatment in the majority of institutions is oral iron supplementation, with blood transfusion reserved for more severe or symptomatic cases.^[3] High doses of oral iron frequently cause side effects, including constipation, nausea and gastric irritation and hence, noncompliance is common. In addition, such therapy has to be given for a long time in cases of severe iron deficiency. Blood transfusions have been used in the treatment of postpartum anaemia, but there are risks associated with its use. Therefore, intravenous iron, alone or in association with recombinant human erythropoietin (rhEPO), has been considered as an alternative in the management of iron deficiency in this setting.

The aim of this study was to find out the prevalence of postpartum anaemia; compare the efficacy of intravenous iron sucrose versus oral iron in subjects with post-partum iron deficiency anaemia, and to study the safety and side effects of these two preparations.

Name & Address of Corresponding Author

Dr. Preeti Sharma
Assistant Professor,
Department of Obstetrics & Gynaecology,
Muzaffarnagar Medical College,
Uttar Pradesh, India.

MATERIALS AND METHODS

The present prospective observational study was conducted in the Department of Obstetrics and Gynecology, Santosh Medical College, Ghaziabad, from January 2013 to December 2013. Total 512 women delivered during this period, out of whom 102 women were detected with postpartum anaemia. Postpartum anaemic women, aged 18 years or more, with diagnosis of iron deficiency anaemia with hemoglobin of 6-9 g/dl, samples taken at 24-48 hours post-delivery and who demonstrated willingness to comply to research protocol were included in the study. Iron deficiency anaemia was diagnosed if hemoglobin was 6-9 g/dl, MCV<78fl, MCH<30pg, serum ferritin <15 microgram/l and microcytic hypochromic to normocytic hypochromic picture on peripheral smear.

Women with hemoglobin>9g/dl, Ferritin >15microgram/l, history of intolerance to iron derivatives, peripartum blood transfusion, history of asthma, thromboembolism, seizures, alcohol or drug abuse, and those having signs of infection or evidence of renal or hepatic dysfunction were excluded from the study.

Written informed consent was taken from all the women, satisfying the inclusion criteria. A detailed history, followed by general physical systemic and local examination was done.

Patients were assigned to two groups of 51 each using computer generated random number sequence. Baseline investigations done on 2nd-3rd day of postpartum included complete blood count with RBC's indices (MCV, MCH MCHC) for severity and type of anaemia, peripheral smear for type of anaemia and serum ferritin for iron stores. There was no statistical significant difference between two groups with respect to these investigations.

One group received 200 mg of oral ferrous sulphate tablets twice daily for six weeks. Other group received intravenous iron sucrose in divided doses on alternate days. Dose was calculated by the following formulae:

$$\text{Weight (in kgs)} \times (\text{Target hemoglobin}-\text{Actual hemoglobin}) \times 0.24 + 500 \text{ mg } 4$$

Target hemoglobin was set at 110 g/L, actual hemoglobin in gram per liter was the patient's hemoglobin level on inclusion; 0.24 was a correction factor and 500 mg for the quantity of stored iron in adults.

Iron sucrose was given in the dose of 100 mg elemental iron, slow intravenous infusions on alternate days (starting from day 2-3 of post delivery). No test dose was required.

Both the groups were supplemented with five milligrams of folic acid.

Result was compared on day 7-14 of delivery (i.e. 1-2 -weeks after treatment) in both the groups. The rate of improvement was measured in terms of hemoglobin, RBC's indices and general

improvement of the patient. At 6 weeks postpartum, the final effect of treatment was judged by repeating all the investigations done as baseline.

Statistical analysis

To compare the difference in means of various parameters (Hb, serum ferritin etc.) between two groups, over period of time (on day 3, 7-14, 42), student-t test was used. P-value < 0.05 was taken as level of statistical significance.

RESULTS

Total 512 women delivered during this period, out of whom 102 women were detected with postpartum anaemia. The prevalence rate of postpartum anaemia in our study was 19.92%.

There was no statistical significant difference between two groups with respect to age of the patients, booked/unbooked/registered, socio-economic status, religion and parity. [Table 1]

Table 1: Socio-demographic factors.

Age (Years)	Sub Group	Oral Group	I/V Group	P-Value
		N=51	N=51	
	19- 24	25	19	0.546
	25-29	19	22	
	30 & above	7	10	
Booking Status	Unbooked	19	25	0.580
	Booked	14	10	
	Registered	18	16	
Religion	Hindu	35	31	0.405
	Muslim	16	20	
Socio-Economic Status	Upper	2	3	0.288
	Middle	10	14	
	Lower	39	34	
Parity	P1	8	9	0.288
	P2	24	15	
	≥P3	19	27	

An increase was seen in the values of hemoglobin, hematocrit, RBC indices and serum ferritin in both the groups. But there was a statistically significant difference between the increase in both the groups, with I.V. group showing much higher values than oral group.

An increase in mean hemoglobin was observed from 7.90± 0.905 g/dl on day 2-3 of postpartum to 8.65± 0.849 g/dl at 1-2 weeks, and to 10.16 ± 0.877 g/dl at 6 weeks postpartum in oral group; whereas from 7.81± 0.849 g/dl on day 2-3 of postpartum to 9.88 ± 0.760 g/dl at 1-2 weeks postpartum and to 10.91 ± 0.770 g/dl at 6 weeks postpartum in I/V group (p value < 0.0001). [Table 2]

Table 2: Comparison of mean hemoglobin between oral and I/V groups.

	On 2nd -3rd day (g/dl)	At 1-2 weeks (g/dl)	At 6 weeks (g/dl)
Oral Group	7.90(±0.905)	8.65 (±0.849)	10.16 (±0.877)
I/V Group	7.81(±0.849)	9.88 (±0.760)	10.91 (±0.77)
p-Value	0.625	0.0001	0.0001

Similarly, there was increase in serum ferritin in both the groups but more in I.V. group. In the oral group, mean serum ferritin on 2nd-3rd postpartum day was 8.04 ± 2.144 ug/L which increased to 23.88 ± 5.339 ug/L at 6 weeks postpartum. In I.V. group, mean serum ferritin was 8.30 ± 1.426 ug/L on 2nd-3rd day postpartum which increased to 58.35 ± 14.537 ug/L at 6 weeks postpartum. This indicates iron sucrose not only treats but also corrects iron stores which is not seen with oral iron. This observation is highly significant in our study thus indicating its use in post-partum anaemia (p value < 0.0001). [Table 3]

Table 3: Comparison of mean serum ferritin between oral and I/V groups.

	On 2nd -3rd day (µg/L)	At 6 weeks (µg/L)
Oral Group	8.04 (± 2.144)	23.88 (± 5.339)
I/V Group	8.30 (± 1.461)	58.35 (± 14.537)
p-VALUE	0.146	0.0001

Comparing the side effects of oral iron and intravenous iron in the present study, in the oral group 20% patients had constipation, 12% had metallic taste, 4% had nausea and vomiting, 2% had abdominal pain and another 2% had diarrhea. The majority of patients (40%) had gastrointestinal side effects. [Figure 1] In I.V. group, 4% patients had mild allergic reaction like rash, flushing and pruritis and 2% had dizziness. 94% patients had no side effects with iron sucrose. In the present study, I.V. sucrose tolerance seems to be excellent and without adverse effects, in accordance with the literature. [Figure 2]




Figure 1: Side effects of oral iron




Figure 2: Side effects of I/V Iron

DISCUSSION

For many pregnant women, postpartum anaemia is inevitable and can be detrimental to the mother and new born. According to the National Pregnancy Nutrition Surveillance System, 29.8% of women who were not previously anaemic during pregnancy become anaemic after delivery.^[5]

Many factors exacerbate postpartum iron status, including maternal characteristics at the start of pregnancy, events occurring during pregnancy, labor and delivery, and the early postpartum period.

The socio-demographic factors in our study were comparable to other studies.

The majority of patients were young and belonged to reproductive age group indicating pregnancy as an important factor for iron deficiency anaemia because of increased demand of iron in pregnancy which is not met by normal diet.^[4-6]

Most of the women belonged to lower socio-economic class taking diet deficient in essential nutrients and minerals and therefore more than 60% women in our study had anaemia. Multiparity had been an important risk factor for postpartum anaemia because repeated pregnancies result in depletion of iron stores.^[6,7]

The majority of patients in all studies were illiterate or educated up to primary school, which could be due to lower socioeconomic status. Due to lack of education, the patients were not aware of the significance of prenatal care and good nutritious diet.^[7]

Women treated with intravenous iron had higher hemoglobin levels in the short term (1-2 weeks postpartum), but no statistical significant difference was observed in patients after 6 weeks in both oral and I.V. iron groups. This indicates the response of iron therapy to be good in both the groups if compliance is assured.^[2,4-9]

Serum ferritin is an indicator of iron stores, and decrease in its values indicates depletion of iron stores. In the present study, there was increase in serum ferritin in both the groups but more in I.V. group. This indicates iron sucrose not only treats but also correct iron stores which is not seen with oral iron. This observation is highly significant in other studies,^[2,4-9] thus indicating its use in postpartum anaemia.

In present study, 40% of patients with oral iron had gastrointestinal side effects, which were comparable with various studies having 30% to 43% of their patients having these side effects. Usually, GI complaints afflict up to 20% of patients taking ferrous iron salts.

In I.V. group, 4% patients had mild allergic reaction like rash, flushing and pruritis and 2% patients had dizziness. 94% patients had no serious side effects with iron sucrose. In the present study, I.V. sucrose tolerance seems to be excellent without adverse

effects, in accordance with literature. These observations were comparable to other studies.^[2,4,6,8] The literature documents that intravenous iron sucrose is reasonably well tolerated (35% of patients have mild side effects) with a low incidence of serious adverse reactions (0.03-0.04%).^[10]

CONCLUSION

Postpartum depression, stress, and lactational failure in women is related to Iron Deficiency Anaemia. These symptoms respond to iron therapy.

Intravenous iron sucrose is an effective alternative to oral ferrous sulphate in the treatment of iron deficiency anaemia during postpartum period. It has shorter treatment periods, a lack of gastrointestinal side effects and produces a more rapid increase in hemoglobin concentration and serum ferritin levels than oral ferrous sulphate. At the same time, it is a safe alternative for the treatment of anaemia, being able to reduce the need for blood transfusion and its concomitant side effect. During postpartum period, it helps to rebuild iron stores, helping the symptoms of anaemia to subside at a faster rate and reduces the risk of developing anaemia in subsequent pregnancies.

Major advantages are safety, efficacy, compliance, simple mode of administration in an outpatient setting and cost effectiveness because admission is not needed in all cases.

Limitations with intravenous iron replacement include the need for medical supervision in the setting of limited health resources.

Acknowledgements

I pay profound regards to my co-faculty and patients in the department of Obstetrics & Gynaecology of Santosh University for their constant support and immeasurable help.

REFERENCES

1. Cochrane Database of Systematic Reviews 2004, Issue 4.
2. Bhandal N, Russell R, Intravenous versus oral iron therapy for postpartum anaemia BJOG 2006; 113: 1248-1252
3. Current Progress in Obstetrics & Gynaecology, John Studd, Volume-1 2012
4. Bayoumeu F, Buisset C., Iron Therapy in iron deficiency anaemia in pregnancy: Intravenous route versus Oral route. Am J Obstet Gynecol 2002; 186 (3): 518 - 522.
5. Pena-Rosas, J.P. Viteri, F.E. (2009) Effects and safety of preventive oral iron or iron+folic acid supplementation for women during pregnancy. Cochrane Database of Systematic Reviews.
6. Momen A.K.A. et al. Intravenous iron sucrose complex in the treatment of iron deficiency anaemia during pregnancy. Eu Journal of Obstet & Gynecol and Reproductive Biology 1996; 69: 121-124.
7. Broche D.E et al. Severe anaemia in the immediate post partum period. Clinical practice and value of intravenous iron. European Journal of Obstetrics & Gynecology and Reproductive Biology 2005; 123: 521-827.
8. Bodnar LM, Cogswell ME. Low income postpartum women are at risk of iron deficiency. J Nutr 2002; 132: 2298 – 302.
9. Perewusnyk G, Breymann C. Parenteral iron therapy in obstetrics: 8 years experience with iron sucrose complex. Br Journal of Nutrition 2002; 88:3-10.
10. Westad S, et al. A 12-week randomized study comparing intravenous iron sucrose versus oral ferrous sulphate for treatment of postpartum anaemia. Acta Obstet Gynecol Scand. 2008; 916-923.

How to cite this article: Sharma P. Prevalence and Management of Postpartum Anaemia in a Tertiary Care Hospital in North India. Ann. Int. Med. Den. Res. 2019; 5(3):OG01-OG04.

Source of Support: Nil, **Conflict of Interest:** None declared