

Staphylococcal Scalded Skin Syndrome in a Child with Nephrotic Syndrome on Steroid Therapy.

Arkya Saha¹, Santosh Kumar²

¹MD student, Department of Pediatric, MGM Medical College & LSK hospital, Kishanganj, Bihar.

²Assistant Professor, Department of Pediatric, MGM Medical College & LSK hospital, Kishanganj, Bihar.

Received: January 2017

Accepted: January 2017

Copyright: © the author(s), publisher. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Staphylococcal scalded skin syndrome (SSSS) is a rapidly progressive exfoliative disease of the skin due to *Staphylococcus aureus* toxins A or B. Prognosis is favourable, the 4% reported deaths occurring mainly because of dehydration or bacterial superinfection.

Keywords: Staphylococcal scalded skin syndrome, nephrotic syndrome, steroid therapy.

INTRODUCTION

Staphylococcal scalded skin syndrome (SSSS) is a rapidly progressive exfoliative disease of the skin due to *Staphylococcus aureus* toxins A or B.^[1] Immunosuppressive states like steroid therapy as in the case of nephrotic syndrome are risk factors for developing such life threatening infections.^[2]

Name & Address of Corresponding Author

Dr. Santosh kumar
Assistant Professor,
Dept of Pediatric,
MGM Medical College & LSK hospital, Kishanganj,
Bihar.

CASE REPORT

A 5-year-old boy, on steroid treatment (prednisolone, 2 mg/kg/day) for the first episode of nephrotic syndrome (3rd week) presented with a generalized vesiculo-bullous rash within 48 hours of evolution of bullous impetigo on the thigh. He had a generalized rash with confluent areas and Nikolsky sign with no mucosal lesions. Throat swab was positive for *Staphylococcus aureus*. Skin biopsy confirmed the diagnosis of SSSS. Steroid was stopped. Treatment with intravenous ceftriaxone and cloxacillin showed good results [Figure 1].

During the treatment course urine protein was negative. Later on the lesions healed without scarring. Since urine protein was negative, 2 weeks later he was started on steroid (1.5 mg/kg alternate day for 6 weeks) to complete the initial treatment of nephrotic syndrome.



Figure 1: Erythematous desquamating lesions after starting treatment (day 4).

DISCUSSION

SSSS is a toxin mediated, epidermolytic condition caused by remote production and haematogenous circulation of toxins that predominantly affect young children. Diagnosis of SSSS is clinical in 70% instances.^[1] Patients with nephrotic syndrome, a common childhood glomerulopathy, are at risk for such life threatening infections, as in this case. First-line treatment comprises anti-staphylococcal penicillins given intravenously along with supportive management. Stress dose is indicated if patient has been given high dose steroids for over a fortnight in the last year.^[2] Prognosis is favourable, the 4% reported deaths occurring mainly because of dehydration or bacterial superinfection.^[1]

REFERENCES

1. Oliveira ARS, Aires S, Faria C, Santos E. Staphylococcal scalded skin syndrome. BMJ Case Reports online: Jun 10, 2013.
Available from: <http://casereports.bmj.com/content/2013/bcr-2013-009478.full.pdf>
2. Bagga A. Revised guidelines for management of steroid-sensitive nephrotic syndrome. Indian Journal of Nephrology 2008; 18:31-9

How to cite this article: Saha A, Kumar S. Staphylococcal Scalded Skin Syndrome in a Child with Nephrotic Syndrome on Steroid Therapy. Ann. Int. Med. Den. Res. 2017; 3(2):PE07-PE08.

Source of Support: Nil, **Conflict of Interest:** None declared