

Study of Conduction Blocks in Acute Myocardial Infarction.

Vijay Kumar¹, Sanjay Goyal², Sanjeev Kumar³, Minal⁴

¹Assistant Professor, Department of Medicine GMC, Patiala.

²Assistant Professor, Department of Medicine GMC, Patiala.

³Professor & Head, Department of Medicine GMC, Patiala.

⁴Junior Resident, Department of Medicine GMC, Patiala.

Received: January 2018

Accepted: January 2018

Copyright: © the author(s), publisher. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Acute Coronary syndromes (ACS) represent a global epidemic. With increase in prevalence of risk factors such as obesity, diabetes and others, there is parallel increase in the incidence of ACS. Which may manifest itself either as acute myocardial infarction (AMI), unstable angina, effort angina or sudden death? Among these the most life threatening is AMI and its associated complications. Conduction blocks occurs in about 15-20% of patients with acute MI. **Methods:** The study was conducted on 100 patients of Acute Myocardial Infarction attending OPD/ admitted in Dept. of Medicine/ ICCU of Rajindra Hospital, Patiala over a period of one Year. Complete clinical examination was done and all the necessary investigations as per case requirement were sent. The patients were observed for conduction blocks for 7 days after the admission or until they stay in the hospital whichever was earlier. **Result:** Out of 100 patients, 21 patients developed conduction blocks. Eight patients (8%) developed first degree AV block, One patient developed mobitz type -2 second degree AV block, 4 patients (4%) developed third degree AV block, 2 patients (2%) developed complete LBBB, 2 Patients (2%) developed LAHB. Overall cardiovascular complications were more in patient with conduction block. The conduction blocks were more common among patients with inferior wall MI than anterior wall MI. The occurrence of cardiogenic shock was more in patients with conduction block. Mortality was higher in patients with block (19.1%) as compared with patients without block (2.5%). **Conclusion:** Conduction blocks are associated with higher in hospital mortality and morbidity in the form of other cardiovascular events during hospital stay, thus conduction blocks are important predictors of poor outcome in patients with AMI. All patients with AMI should be watched carefully for early recognition of conduction blocks and appropriate treatment should be started early.

Keywords: Conduction Block, Acute myocardial Infarction.

INTRODUCTION

Acute Coronary syndrome (ACS) represents a Global epidemic. With increase in prevalence of risk factors such as obesity, diabetes and others, there is parallel increase in the incidence of ACS.

ACS may manifest itself either as acute myocardial infarction (AMI), unstable angina, effort angina or sudden death. Among these the most life threatening is AMI and its associated complications.

Conduction blocks occur in about 15-20% of patients with acute MI, studies have shown that conduction blocks are important predictors of poor outcome in patients with acute MI and are associated with higher in hospital mortality and other cardiovascular events in the form of hypotension

requiring treatment, left ventricular failure, cardiogenic shock, recurrent angina, complete AV block and cardiac arrest.^[1] In this study an attempt is made to know the association of conduction blocks and their prognostic value in patients of acute MI.

Aims & Objectives

1. To study the incidence and type of conduction blocks in Acute Myocardial Infarction.
2. To study the prognostic significance of various conduction blocks in Acute Myocardial Infarction.

MATERIALS AND METHODS

The study was conducted on 100 patients of Acute Myocardial Infarction attending OPD/ admitted in Dept. of Medicine/ ICCU of Rajindra Hospital, Patiala over a period of one year. Written consent for the study was obtained from all patients after examining them in detail and consent of institutional, ethical committee was obtained.

Inclusion criteria

- Patients diagnosed as having acute myocardial infarction on clinical and ECG or cardiac markers basis.

Name & Address of Corresponding Author

Dr. Sanjay Goyal
Assistant Professor,
Department of Medicine
Govt Medical College,
Patiala, Punjab,
India.

- Diagnosed as having conduction blocks based on ECG during first seven days of acute myocardial infarction.

Exclusion Criteria

- Previously known cases of conduction blocks.
- Patients already on drugs which may cause conduction blocks, like Beta blockers, calcium channel blockers and Digoxin.
- Inability to get informed consent.

A detailed history of each patient was taken as per proforma attached. Complete clinical examination was done and all the necessary investigations as per case requirement were sent. The patients were observed for conduction blocks for 7 days after the admission or until they stay in the hospital whichever was earlier. The diagnosis of various conduction blocks like Left bundle branch block,^[15] Right bundle branch block,^[15] Left anterior hemiblock,^[15] Left posterior hemiblock,^[15] Bifascicular blocks,^[17] Trifascicular block,^[17] First

degree & Second degree AV Block were made based on the ECG features.^[16]

Criteria for Acute Myocardial Infarction

Either one of the following criteria satisfies the diagnosis for an acute, evolving MI:

- Ischaemic symptoms;
- ECG changes indicative of ST elevation MI or non ST elevation MI.
- Rise in cardiac biomarkers such as CPK-MB or Troponin-T or both.

Statistical tests:

The data collected were analyzed using proper tests:

- Student t test
- Chi-square test for association

Continuous variables are presented as mean ± SD and frequency variables as percentages. P value of <0.05 was considered for statistical significance.

RESULTS

Table 1: Age Distribution of Patients According To Conduction Blocks

| Age Groups (in years) | Conduction Block | | | | Total |
|-----------------------|------------------|------|---------|------|-------|
| | Absent | | Present | | |
| | No. | % | No. | % | |
| <30 | 1 | 1.3 | 0 | 0 | 1 |
| 31-40 | 7 | 8.9 | 2 | 9.5 | 9 |
| 41-50 | 16 | 20.3 | 2 | 9.5 | 18 |
| 51-60 | 27 | 34.2 | 6 | 28.6 | 33 |
| 61-70 | 18 | 22.8 | 7 | 33.3 | 25 |
| 71-80 | 10 | 12.7 | 3 | 14.3 | 13 |
| > 80 | 0 | 0 | 1 | 4.8 | 1 |
| Total | 79 | 100 | 21 | 100 | 100 |

Table 2: Gender Distribution of Patients According To Conduction Block.

| Gender | Conduction Block | | | | Total |
|--------|------------------|------|---------|------|-------|
| | Absent | | Present | | |
| | No. | % | No. | % | |
| Male | 62 | 78.5 | 13 | 61.9 | 75 |
| Female | 17 | 21.5 | 8 | 38.1 | 25 |
| Total | 79 | 100 | 21 | 100 | 100 |

Table 3: Various Sites Of AMI

| Site | No. of cases | Percentage |
|---------------------|--------------|------------|
| Anterior | 37 | 37 |
| Antero lateral | 12 | 12 |
| Antero septal | 13 | 13 |
| Inferior | 37 | 37 |
| Inferior with right | 1 | 1 |
| Total | 100 | 100 |

Table 4:- Incidence Of Various Types Of Conduction Blocks In The Present Study (N=100)

| Types of conduction block | No. of cases | Percentage |
|---------------------------|--------------|------------|
| First degree AV block | 8 | 8 |
| Second degree AV block | 1 | 1 |
| Third degree AV block | 4 | 4 |
| LBBB | 2 | 2 |
| RBBB | 4 | 4 |
| LAHB | 2 | 2 |
| Total | 21 | 21 |

Table 5: Association Of Site Of Mi With Conduction Blocks.

| Site of MI | Conduction Block | | | | | | | Total |
|-------------|------------------|------------|-------------------|------------|--------|--------|--------|-------|
| | Absent | 1st degree | 2:1 Mobitz type 2 | 3rd Degree | LBBB | RBBB | LAHB | |
| INF with RT | 1(1.3) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 1 |
| ANT | 32 (40.5) | 1(12.5) | 0 (0) | 1(25) | 2(100) | 0 (0) | 1(50) | 37 |
| ANT-LAT | 11(13.9) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 1(25) | 0 (0) | 12 |
| ANT-SEP | 12(15.2) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 1(50) | 13 |
| INF | 23(29.1) | 7(87.5) | 1(100) | 3(75) | 0 (0) | 3(75) | 0 (0) | 37 |
| Total | 79 (100) | 8(100) | 1(100) | 4(100) | 2(100) | 4(100) | 2(100) | 100 |

Table 6:- Association Of Conduction Blocks With Complications

| Site of MI | Absent | 1st degree | 2:1 Mobitz type 2 | 3rd Degree | LBBB | RBBB | LAHB | Total |
|-------------------|----------|------------|-------------------|------------|--------|--------|--------|-------|
| Absent | 64(81.1) | 5(62.5) | 1 (100) | 1 (25) | 1 (50) | 1 (25) | 1 (50) | 74 |
| CHF | 5 (6.3) | 1(12.5) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 6 |
| Death | 2 (2.5) | 0 (0) | 0 (0) | 2 (50) | 0 (0) | 2 (50) | 0 (0) | 6 |
| LVF | 5(6.3) | 0 (0) | 0 (0) | 0 (0) | 1(50) | 0 (0) | 0 (0) | 6 |
| Cardiogenic Shock | 2 (2.5) | 2 (2.5) | 0 (0) | 1 (25) | 0 (0) | 1 (25) | 1 (50) | 7 |
| VT | 1 (1.3) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 1 |
| Total | 79(100) | 8 (100) | 1(100) | 4(100) | 2(100) | 4(100) | 2(100) | 100 |

Table 7:- Mortality In Patients With Conduction Blocks.

| Outcome | Without conduction block n=79 | With Conduction Block n=21 | | | | | | Total N=100 |
|---------|-------------------------------|----------------------------|-------------------|------------|----------|----------|----------|-------------|
| | | 1st degree | 2:1 Mobitz type 2 | 3rd Degree | LBBB N=2 | RBBB N=4 | LAHB N=2 | |
| Deaths | 2(2.5) | 0 (0) | 0 (0) | 2 (50) | 0 (0) | 2 (50) | 0 (0) | 6 (6) |

The present study was a hospital bound study. 100 patients of Acute Myocardial Infarction were included considering inclusion and exclusion criteria. Of them 21 patients developed conduction blocks.

Mortality was higher in patients with blocks (19.1%) as compared to patients without blocks (2.5%). It was significant statistically with p value 0.008.

DISCUSSION

Conduction blocks are frequent complications of acute myocardial infarction. This study has documented the frequency of conduction blocks and

their influence on prognosis in 100 patients with acute myocardial infarction. Blocks at the atrioventricular node occurred almost exclusively in inferior infarction, while blocks involving the bundle branches were more common in anterior infarction.

In the present study, the mean age of patient with conduction blocks was 62.90 whereas in patients without blocks was 57.37. On comparison with other studies, our figures are in concordance with the study done by Newby KH et al and Abidov et al.^[10,13] In the present study group 75% were males and 25% were females. On doing comparison with other studies, our figures are in concordance with the study done by Newby KH et al.

Table 8: Comparison Of Incidence Of Av Blocks In Present Study With Previous Studies.

| | Present Study | Meltzer & Kitchell ^[14] | Hurwitz & Eliot |
|------------------------|---------------|------------------------------------|-----------------|
| First degree AV Block | 8% | 8.5% | 5.8% |
| Second degree AV Block | 2% | 3.5% | 2% |
| Third degree AV Block | 4% | 4.2% | 1.8% |

Table 9: Comparison Of Incidence Of Rbbb And Lbbb In The Present Study With The Previous Studies.

| | Present Study | Stephen scheidt & Thomas Killip ^[3] | Col & Weinberg ^[4] | Rizzon, Biase & Baissus ^[5] |
|------|---------------|--|-------------------------------|--|
| RBBB | 4% | 3% | 3.3% | 3% |
| LBBB | 2% | 3% | 3.7% | 2% |

In the present study the incidence of RBBB was more than LBBB. It was comparable with the results

of Stephen scheidt & Thomas Killip,^[3] Col & Weinberg and Rizzon, Biase & Baissus.^[4,5] Stephen

scheidt & Thomas Killip have reported an equal incidence of RBBB and LBBB.^[3] However some studies have noted a higher incidence of RBBB than LBBB. Two patients with RBBB expired with mortality of 50%. Two patients (2%) in the present study developed left anterior hemiblock (LAHB), which is comparable with 4.7% reported by James atkins et al.^[8]

None of the patients in present study developed left posterior hemiblock (LPHB). It is usually rare. An incidence of 0.3% was reported by Rizzon, Biase & Baissus.^[5] The conduction blocks were significantly more common among patients with inferior wall MI (66.7%) than the anterior wall MI (33.3%) with a p value of 0.016 which was statistically significant. The results are complete with study of Majumdar AA et al,^[11] which also showed higher incidence of conduction blocks in inferior wall MI patients than anterior myocardial infarction (56.8% and 31.8% respectively).

Conduction blocks were mostly atrioventricular (78.6%) in inferior wall MI where as they were mostly intraventricular (71.4%) with anterior wall MI. These results are complete with the study done by Majumdar AA et al,^[11] which are 92% and 72% respectively. In our study the mortality rate in patients with complete AV block was 50%, which was higher than that of patients without blocks which was 2.5% and this is comparable to study done by Goldberg et al,^[2] but is higher than study by Beher et al.^[9]

Among the 4 patients who developed RBBB in the present study, 2 patients expired with a mortality of 50%, which is in concordance with 52% as in Godman, Lassers & Julian^[7], but is higher than the results of Moreno AM et al.^[12]

Our values are comparatively lesser when compared to study done by Majumdar AA et al.^[11]

Table 10: Comparison Of Overall Mortality In Patients With Blocks In Our Study And Other Studies.

| Studies | Mortality rate (%) | |
|-------------------|------------------------------------|---------------------------------|
| | Without CBs (Complete Heart Block) | With CBs (Complete Heart Block) |
| Present study | 2.5 | 19.1 |
| Majumdar AA et al | 3.6 | 25 |

Table 11: Comparison Of Frequency Of Cardiogenic Shock In Patients With Iii Degree Block In Our Study And Other Studies.

| Studies | With III Block (%) | Without Block (%) |
|------------------------------------|--------------------|-------------------|
| Present Study | 25 | 2.5 |
| Beher S. et al ^[9] | 23 | 4 |
| Goldberger RJ et al ^[2] | 23.4 | 6.5 |

Table 12: Comparison Of Frequency Of Cardiogenic Shock In Patients With Rbbb Block In Our Study And Other Study.

| Studies | With RBBB (%) | Without RBBB (%) |
|----------------------------|---------------|------------------|
| Present Study | 25 | 2.5 |
| GO AS et al ^[5] | 10 | 5.6 |

Our values are similar with other studies. All studies showed higher incidence of cardiogenic shock with complete heart block.

In RBBB patients 25% developed cardiogenic shock as compared to 2.5% in patients without block on comparison with other studies.

Our values are higher when compared with study done by GO AS et al.^[5]

CONCLUSION

The most common conduction block in the present study was first degree AV block (8%), followed by Right Bundle Branch Block (RBBB) (4%), third degree or complete AV block (4%), Left Anterior Hemiblock (LAHB) (2%), Left Bundle Branch Block (LBBB) (2%), second degree AV block (1%). All patients with second degree AV block in the present study, had Mobitz type 2 variety and all patients with RBBB and LBBB had complete RBBB and complete LBBB respectively. Among 13

patients with anteroseptal MI, 1 patient (7.7%) developed left anterior hemiblock.

The conduction blocks were significantly more common among patients with inferior wall MI, than the anterior wall MI. Bundle branch blocks were more common in anterior wall MI than inferior wall MI; whereas antriventricular blocks were more common in inferior wall MI. Bradycardia, Hypotension and raised JVP were more common in patients with conduction blocks as compared to patients without conduction blocks. Most common complications were CCF, LVF and cardiogenic shock among both groups. These complications were seen more commonly in patients with blocks. Mortality was higher in patients with blocks (19.1%), as compared to patients without blocks (2.5%), various pattern of conduction blocks developed following AMI and they have a varied impact on the outcome, conduction blocks are associated with higher in hospital mortality and morbidity in the form of other cardiovascular events

during hospital stay, thus conduction blocks are important predictors of poor outcome in patients with AMI. All patients with AMI should be watched carefully for early recognition of conduction blocks and appropriate treatment should be started early.

How to cite this article: Kumar V, Goyal S, Kumar S, Mirnal. Study of Conduction Blocks in Acute Myocardial Infarction. *Ann. Int. Med. Den. Res.* 2018; 4(2):ME20-ME24.

Source of Support: Nil, **Conflict of Interest:** None declared

REFERENCES

1. Go AS, Barron HV, Rundle AC, Ornato JP, Avins AL. Bundle branch block and in hospital mortality in Acute Myocardial Infarction. *ANN Intern Med.*
2. Goldberg RJ, Zevallos JC, Yarzebski J, Alpert JS, Gore JM, Chenz, et al. Prognosis of acute myocardial infarction complicated by complete heart block (The Worcester Heart Attack study). *Am J Cardiol* 1992;69: 1135-41.
3. Scheidt S, Killip T. Bundle branch block complicating acute myocardial infarction. *JAMA* 1972; 222(8): 919-24.
4. Col JJ, Weinberg SL. The incidence and mortality of intraventricular conduction defects in acute myocardial infarction. *Am J Cardiol* 1972;29: 344-50.
5. Rizzon P, Biase MD, Baissus C. Intraventricular conduction defects in acute myocardial infarction. *Br Heart J* 1974;36: 660-68.
6. Milton H, Robert S. Eliot. Arrhythmias in Acute Myocardial Infarction. *Chest.* 1964;45(6): 616-26.
7. Godman M J, Lassers B W, Julian D G. Complete Bundle Branch Block Complicating Acute Myocardial Infarction. *N Engl J Med* 1970; 282: 237-40.
8. Atkins JM, Leshin SJ, Blomqvist G, Mullins CB. Ventricular conduction blocks and sudden death in acute myocardial infarction. *New Engl J Med* 1973; 288(6): 281-284.
9. Beher S, Zissman E, Zion M, Shalev Y, Capsi A, Kaplinsky et al. Complete Atrioventricular block complicating inferior wall acute MI: Short and long term prognosis. *Am Heart J* 1993; 125:1622-6.
10. Newby KH, Pisano E, Krucoff MW, Green C, Netale A. Incidence and clinical relevance of the occurrence of bundle branch block in patients treated with thrombolytic therapy. *Circulation* 1996;94: 2424-8.
11. Majumder AA, Malik A, Zafar A. Conduction disturbances in acute myocardial infarction: incidence, site wise relationship and the influence of in-hospital prognosis. *Bangladesh Med Res Counc Bull* 1996;22 (2): 74-80.
12. Moreno AM, Thomas GJ, Alberola GA, Chavarri MV, Soria FC, Sanchez EM et al. Incidence, clinical characteristics and prognostic significance of Right Bundle branch block in Acute MI: A study in Thrombolytic era *Circulation* 1997; 96: 1139-44.
13. Abidov A, Kaluski E, Hod H, Leor J, Vered Z, Becher G et al. Influence of conduction disturbance on clinical outcome in patients with Acute Myocardial infarction receiving thrombolysis (results from ARGAMI -2 study). *Am J Cardiol* 2004;93: 76-80.
14. Meltzer LE, Kitchell JB. The incidence of arrhythmias associated with acute myocardial infarction. *Progress in Cardiovascular Disease* 1966;9 (1): 50-63.
15. Schamroth L. An introduction to Electrocardiography. 7th ed. Oxford: Black Well science 1990; 89-114.
16. Goldberger AL. Clinical Electrocardiography: A simplified approach. 8th ed. Saunders: Elsevier 1999; 159-162.
17. Goldberger AL. Clinical Electrocardiography: A simplified approach. 8th ed. Saunders: Elsevier 1999; 62-63.