

Antibiotics Prescribing Pattern in Pediatrics Ward in Tertiary Care Hospital.

Meherunisa¹, Sapna Jaiswal², Zaigam Abbas³

¹Associate Professor, Department of Pharmacology, Carrier Institute of Medical Sciences, Lucknow.

²Tutor, Department of Biochemistry, Ram ManoharLohia Institute of Medical Sciences, Gomti Nagar, Lucknow, Uttar Pradesh, India.

³Associate Professor, Department of Pedia, Carrier Institute of Medical Sciences, Lucknow.

Received: September 2018

Accepted: September 2018

Copyright:© the author(s), publisher. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Aims & Objective:- The present study was suggested with the objective to of antimicrobial drug use in different indication in the tertiary care hospital. **Methods:** The case study was carried out in the Departments of pharmacology and pediatrics of carrier institute of Medical sciences Lucknow. **Results:** In the present study, total 250 cases were included. Among the 250 cases 70% were male and 30% were female. In this study we found that 49.2% cases were belong to between 1year to 5 year age group followed by 1 month – 1 year (14.8%) &>5 year (36%).In our study , 43.2% cases had 6-12 kg weight followed by 28%(12-18 kg) , 16.8%(18-30 kg), 12%(<6kg). **Conclusion:** This study conclude that, common use of antibiotic without culture and sensitivity, may leads to high cost burden and commonly resistance to antibiotics. This situation comes from parents' pressure of quick relief using strong medication.

Keywords: Children, pharmacology, antimicrobial drug, cephalosporin's

INTRODUCTION

Fever is one of most common clinical symptoms managed by pediatricians and other health care providers and accounts up to one third of all presenting conditions in children.^[1] Fever is not an illness but a physiologic mechanism that has beneficial effects against infection.^[2-4] Fever primarily enhances neutrophil production and T-lymphocyte proliferation, retards the growth and reproduction of bacteria and viruses, and aids in the body's acute-phase reaction.^[5-9] Data show beneficial effects on certain components of the immune system in fever, and limited data have revealed that fever actually helps the body recover more quickly from viral infections, although the fever may result in discomfort in children.^[5,10-12]

Children constitute about 40% of India's population. Infants and children suffer from frequent but usually non serious illnesses. Most of these are self-limiting and are often treated not only inappropriately, but also resorting to poly-pharmacy.^[13,14] The key role of antibiotics for the treatment of infectious diseases that are prevalent everywhere in developing

countries may not be denied. However, there are also reports of an irrational use of antibiotics which may even lead to infections that are worse than the originally diagnosed ones.^[15,16]

MATERIALS AND METHODS

Study population

250 cases were included in this study.

Study Area

The case study was carried out in the Departments of pharmacology and pediatrics of carrier institute of Medical sciences Lucknow.

Study duration

Duration of this study was one year.

Sampling technique & Data collection

Study population included of patients who were admitted in pediatrics who had with primary complain of fever. Therefore, all of the cases admitted on the data collection period were covered and it came to by complete enumeration method. Patient related information such as age, sex, diagnosis and drug related information like drugs, dose, dosage form, route of administration were collected in a pre-designed, structured schedule in accordance with WHO criteria for Drug Utilization Survey and Research.

Data Analysis

Data were analyzed by using Microsoft excel.

Name & Address of Corresponding Author

Dr. Meherunisa
Associate Professor,
Department of Pharmacology,
Carrier Institute of Medical Sciences,
Lucknow.

RESULTS

Table 1: Gender Wise Distribution

Gender	Number Of Cases	Percentage
Male	175	70%
Female	75	30%
Total	250	100%




Figure 1: This chart shows distribution of cases in males & females

Table 2: Age Wise Distribution.

Age	Number Of Cases	Percentage
1 M – 1 Y	37	14.8%
1Y – 5 Y	123	49.2%
>5 Y	90	36%
TOTAL	250	100%




Figure 2: This chart shows distribution of cases accordingly age in all cases

Table 3: Weight Wise Distribution

Weight	Number Of Cases	Percentage
<6	30	12%
6-12	108	43.2%
12-18	70	28%
18-30	42	16.8%
TOTAL	250	100%

In the present study, total 250 cases were included. Among the 250 cases 70% were male and 30% were female. In this study we found that 49.2% cases were belong to between 1year to 5 year age group followed by 1 month – 1 year (14.8%) & >5 year (36%). In our study, 43.2% cases had 6-12 kg weight followed by 28% (12-18 kg), 16.8%(18-30 kg),

12%(<6kg). Among the all cases we found that 24.8% diagnosis had of gastrointestinal tract infection followed by LRTI (22.4%) ,labor pneumonia (1.2%), bronchopneumonia(0.8%), emphyma (0.8%), UTI(16.4%), viral fever(21.2%), unknown fever (12.4%). In the all cases, most prescribing the drug were from cephalosporin group (40.8%) followed by penicillin(21.2%), aminoglycoside (27.2%) and other group (10.8%).




Figure 3: This chart shows distribution of cases according to weight

Table 4: Diagnosis in all cases

Diagnosis	Number Of Cases	Percentage
Lower Respiratory Tract Infection	56	22.4%
Labor Pneumonia	3	1.2%
Bronchopneumonia	2	0.8%
Emphyma	2	0.8%
Gastrointestinal Infection	62	24.8%
Urinary Tract Infection	41	16.4%
Viral Fever	53	21.2%
Uknown Fever	31	12.4%
Total	250	100%

Table 5: WHO recommended Prescribing Indicators

Average number of drugs prescribed per day	3.21%
Percentage of the drugs prescribed by generic name	24.39%
Percentage of the drugs prescribed from essential drug list	96.58%
Percentage of the drugs supplied from hospital pharmacy	65.71%
Percentage of inject able drugs prescribed per day	67.60%
Percentage of prescription containing antimicrobial agent	94%

Table 6: Precribing All Group Of Drug In All Cases

Group Of Drug	Number Of Cases	Percentage
Penicillin	53	21.2%
Cephalosporin	102	40.8%
Aminoglycoside	68	27.2%
Other	27	10.8%
Total	250	100%

Table 7: Prescribing Penicillin Group Of Drug In All Cases

Penicillin Group	Number Of Cases	Percentage
Penicillin	1	1.9%
Ampicillin	8	15.1%
Amoxicillin	9	16.9%
Amoxicillin+clavulanic acid	31	58.4%
Piperacillin + tazobactam	4	7.5%
TOTAL	53	100%

Table 8: Prescribing Cephalosporin Group Of Drug In All Cases

Cephalosporin Group	Number Of Cases	Percentage
Ceftriaxone	38	37.2%
Ceftriaxone + Sulbactam	14	13.7%
Ceftriaxone + Tazobactam	7	6.8%
Cefotaxim	19	18.6%
Cefepime	16	15.6%
Cefpodoxim	2	1.9%
Ceftazidime	6	5.8%
TOTAL	102	100%

Table 9: Prescribing Aminoglycoside Group Of Drug In All Cases

Aminoglycoside group	Number of cases	Percentage
Amikacin	12	17.6%
Gentamycin	49	72.1%
Tobramycin	7	10.3%
TOTAL	68	100%

DISCUSSION

In our study, was noted to be high in male cases than the female and in patients of 1 year to 5 years of age. These might be related to prevalence of infectious diseases in these segments of pediatric populations. Antibiotic use is seemed to be indiscriminate as 98.8% of patients admitted with fever got at least one antimicrobial drug whereas documented diagnoses revealed that in 21.2% cases the cause was viral. Some of the patients who received antimicrobial drugs, take three or more antibiotics simultaneously which might be predispose to higher chance of antibiotic resistance. Moreover in case of prescriptions of UTI, culture and sensitivity reports were not documented which indicate those patients were treated empirically with antimicrobial drugs.

WHO define that the rational use of drugs when “patients receive medications appropriate to their clinical needs, in doses that meet their own individual requirements, for an adequate period of time, and at the lowest cost to them and their community”.^[17] The irrational use of drugs is known to lead to an increase in the cost of treatment, incidence of ADR and development of resistance against antimicrobials.^[18] There is enough evidence to demonstrate that prescribing of drugs has shifted

from generics to branded and prescribing out of NLEM.^[19-21]

The results of studies found to help policy makers to develop policy regarding quality of rational drug use in a health facility.^[22-24]

CONCLUSION

Frequent use of antibiotic without frequent culture sensitivity, leads to high cost burden and resistance to antibiotics. This situation comes from parents' pressure of quick relief using strong medication.

REFERENCES

- Crocetti M, Moghbeli N, Serwint J. Fever phobia revisited: have parental misconceptions about fever changed in 20 years. *Pediatrics*. 2001; 107(6):1241–1246
- Bilenko N, Tessler H, Okbe R, Press J, Gorodischer R. Determinants of antipyretic misuse in children up to 5 years of age: a cross-sectional study. *Clin Ther*. 2006;28(5): 783–793
- Kramer MS, Naimark L, Leduc DG. Parental fever phobia and its correlates. *Pediatrics*. 1985;75(6):1110–1113
- Li SF, Lacher B, Crain EF. Acetaminophen and ibuprofen dosing by parents. *Pediatr Emerg Care*. 2000;16(6):394–397
- Jaffe DM. Assessment of the child with fever. In: Rudolph CD, Rudolph AM, Hostetter MK, Lister GE, Siegel NJ, eds. *Rudolph's Pediatrics*. 21st ed. New York, NY: McGraw- Hill; 2002:302–309
- Kohl KS, Marcy SM, Blum M, et al; Brighton Collaboration Fever Working Group. Fever after immunization: current concepts and improved future scientific understanding. *Clin Infect Dis*. 2004;39(3):389–394
- Hasday JD, Garrison A. Antipyretic therapy in patients with sepsis. *Clin Infect Dis*. 2000; 31(suppl 5):S234–S241
- Adam HM. Fever and host responses. *Pediatr Rev*. 1996;17(9):330–331
- Kluger MJ. Fever revisited. *Pediatrics*. 1992; 90(6):846–850
- Roberts NJ. Impact of temperature elevation on immunologic defenses. *Rev Infect Dis*. 1991;13(3):462–272
- Nizet V, Vinci RJ, Lovejoy FH. Fever in children. *Pediatr Rev*. 1994;15(4):127–135
- Kluger MJ, Roberts NJ. Fever. *Pediatrics*. 1990; 96(6):886–890
- Straand J, Rokstad K, Heggedal U. Drug prescribing. *Acta Paediatr*. 1998;(87): p. 218-24.
- Ghai OP, Paul VK. Rational drug therapy in pediatric practice. *Indian Pediatr*. 1988; 25:p. 1095-109.
- Principi N. Control of antibiotic therapy in paediatric patients. *Developmental pharmacology and therapeutics*. 1981; 2: p. 145-55.
- Schollenberg E, Albritton WL. Antibiotic misuse in a paediatric teaching hospital. *Can Med Assoc J*. 1980;122: p. 49-52.
- Department Of Health And Family Welfare, Government Of West Bengal. *Standard Treatment Guideline For Primary health Care Facilities*. 2nd ed. Kolkata; 2011.
- World Health Organization. *Recommendations for management of common childhood conditions Switzerland*; 2012.
- Rational use of medicine. World Health Organisation site.[Online]. [cited 2011 Dec 2]. Available from: http://www.who.int/medicines/areas/rational_use/en/.
- Cheraghali AM, Idries AM. Availability, affordability, and prescribing pattern of medicines in Sudan. *Pharm World Sci*. 2009; 31(2): p. 209-15.
- Drug utilization. (IUPHAR). [Online]. [cited 2011 Dec 2]. Available from: http://www.iuphar.org/pdf/hum_76.pdf.

22. Adebayo ET, Hussain NA. Pattern of prescription drug use in Nigerian army Hospitals. *Ann Afr Med.* 2010; 9(3): p. 152-8.
23. Jhaj R, Bhargava VK, Uppal R, Reeta K, Saha L, Kaur N, et al. Drug prescribing in children in a North Indian referral hospital. *Pharmacoepidemiol Drug Saf.* 2000; 9(5): p. 423-7
24. Bimo H, Hogerzeil V. How to Investigate Drug Use in Health Facilities- Selective Drug Use Indicator. In WHO Department of Essential Drug and Medicine policy 1999.p. 3.

How to cite this article: Meherunisa, Jaiswal S, Abbas Z. Antibiotics Prescribing Pattern in Pediatrics Ward in Tertiary Care Hospital. *Ann. Int. Med. Den. Res.* 2018; 4(6):PC01-PC04.

Source of Support: Nil, **Conflict of Interest:** None declared