

# Exercise prescriptions in Obesity, Diabetes and Hypothyroidism: A Review.

Nitin John<sup>1</sup>, Jyoti John<sup>2</sup>

<sup>1</sup>Professor and Head, Department of Physiology, Dr Ram Manohar Lohia Institute of Medical Sciences, Lucknow, Uttar Pradesh, India.

<sup>2</sup>Associate Professor, Department of Biochemistry, Dr Ram Manohar Lohia Institute of Medical Sciences, Lucknow, Uttar Pradesh, India.

Received: February 2018

Accepted: February 2018

**Copyright:** © the author(s), publisher. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

Exercise is being recommended to patients by Physician and Exercise Physiologist over the years. Many research studies conducted in last decade pointed out the benefits of exercise on health. Various researchers conducted different exercise protocol to evaluate its effect on health. By and large most patients or individuals with coronary vascular risk factors, or those suffering with type 2 diabetes, Obesity induced hypertension, metabolic syndrome, or hypothyroid obese practice their self-design exercise regime or one recommended by any health gymnasium instructor. But as Obesity, Diabetes and Metabolic Syndrome have reached the iceberg and are main contributors towards mortality and death; it's necessary that apart from dietary control a well-designed scientific exercise protocol should be followed by the diseases. These facts gave us an impetus to earmark suitable exercise and prescription regime in Obese, Diabetics and Hypothyroid Obese.

**Keywords:** Obesity, Hypertension, Diabetes, Hypothyroid, Physician, Exercise Physiologist.

## INTRODUCTION

Exercise and its benefits are widely acclaimed by doctors, researchers and scientist community. The benefits of exercise are immense and it promotes longevity of life; helps in weight control, aids in socialization, and help in prevention of diseases. Exercise promotes optimum health. As a matter of fact most healthy individuals and patients especially of obesity, diabetes and hypertension practice some kind of physical activity or exercise protocol which is either self-determined or as instructed by health gym trainer. But practicing a scientifically based health prescription is the need of the day today. All individual's healthy and diseased needs to consult Physician and Exercise Physiologist for exercise prescription. Patients with cardiorespiratory diseases, metabolic syndromes, arrhythmia, acute pulmonary embolism, acute myocarditis, sickle cell anemia, thalassemia etc. needs special attention.<sup>[1,2]</sup> These facts gave us an impetus to explore the literature for finding out the various exercise prescription regime in Obesity, Diabetes and Hypothyroidism.

### Name & Address of Corresponding Author

Dr Nitin John,  
Professor and Head,  
Department of Physiology,  
Dr Ram Manohar Lohia Institute of Medical Sciences,  
Lucknow,  
Uttar Pradesh.

## DISCUSSION

Overweight and obesity varied considerably across the globe. Overweight is found to be in 9% of population in India while it is 37% in Russia. The percentage of obese is around 3% in India to about 35% in South Africa.<sup>[2]</sup> The prevalence of hypertension, angina, stroke and diabetes is considerably at rise in obese and overweight individuals. It is very much needed that the exercise which has proven to benefit the obese individuals in controlling weight and occurrence of hypertension, diabetes and hypothyroidism needs to be inculcated as way of life in overweight and obese individuals to prevent morbidity and mortality in them.

The benefits of exercise on physiological functions and biochemical homeostasis; as revealed by various scientific study needs to be emphasized and discussed.

### Exercise and Calcium Homeostasis

Physical exercise improves bone mass predominantly at the load-bearing bone sites. Insulin-like growth factor-1 and sexual hormones which are altered by physical training exercises modulates the bone response. Maimoon L et al found that exercise influences calcium homeostasis and alter calciotropic hormone levels depending on duration and intensity of exercise training.<sup>[3]</sup> Henderson SA et al found out that serum calcium level is significantly increased by exercise at 50% of maximum aerobic capacity (VO<sub>2</sub> max), and

this persists until the late recovery phase and thereafter it falls below resting levels. The plasma parathormone levels are initially decreased by moderate exercise (50% VO<sub>2</sub> max.) but get significantly increased during and immediately after severe exercise. Exercise stimulates the hormonal, osteolytic influence on the skeleton, increases plasma parathormone release and mobilises calcium stores producing a positive skeletal balance.<sup>[4,8]</sup>

### **Exercise and Diabetes Prevention**

Exercise is the best insulin sensitizer and its benefits are immense. The physiological understanding of exercise explains that during the first few minutes of exercise the muscle glycogen breaks over anaerobically. It's after 5-10 minutes of exercise the muscle glycogenolysis activity slower down and then the glucose from the liver is released into circulation and is taken up by the muscles as fuel by process hepatic glycogenolysis. After twenty minutes of completing the exercise the muscles' glycogen stores are now depleted and the blood glucose is now maintained by hepatic glycogenolysis and triglycerides that are broken down from adipose tissue by lipolysis. The further continuation of exercise leads to breaking down of fat to free fatty acids (FFA) and is used as a source of fuel for the muscles. The benefits of exercise in diabetes is that it revitalize blood glucose control, improves insulin sensitivity, lowers insulin requirements often leading to a reduced dosage of insulin and or oral hypoglycemic agents especially in patients of Type 2 diabetes. In the earlier phase of exercise there is translocation of glucose transporters to the plasma membrane and glucose uptake. As the exercise duration progresses there is up regulation of glucose transporter numbers and changes in capillary density and this causes an improvement in insulin sensitivity. Exercise is beneficial for both glucose uptake mechanisms and the anti-lipolytic effects of insulin.<sup>9</sup> Physical activity and exercise training also benefit patients with type 1 diabetes mellitus (T1DM) reducing the risk of cardiovascular diseases and metabolic syndrome. T1DM patients avoid performing exercise due to the apprehension of developing hypoglycemia and this can be prevented by modulating the dose of both basal long-acting insulin and short-acting insulin, as well as the choice of exercise intensity and duration.<sup>[10]</sup>

### **Exercise and Brain Region Specific Adaptation**

Julie A Morgan et al and Greenwood BN et al pointed out in their review as they investigated regional brain adaptations to exercise, that there is evidence of multiple regional adaptations to both forced and voluntary exercise.<sup>[11]</sup> Exercise can induce molecular adaptations in neuronal functions in several instances.<sup>[12-15]</sup> Similarly Mischel et al found out that physical (in) activity-dependent structural plasticity in bulbospinal catecholaminergic

neurons of rat rostral ventrolateral medulla.<sup>[16]</sup> The physiological adaptations in exercises may elucidate molecular and cellular mechanisms of recovery in psychiatric and neurological health conditions.

### **Zinc Metabolism in Exercise**

Exercise influences zinc metabolism and there redistribution within the body. There is increased zinc requirement with strenuous activity due to additional zinc losses via sweat and urine as evaluated by Overgaard K et al, DeRuisseau KC et al in their studies.<sup>[17-19]</sup>

### **Effects of training on potassium homeostasis during exercise**

Potassium which is released by contracting skeletal muscle cells facilitates muscle contraction but it also leads to muscular fatigue. Endurance and sprint training prolonged and high intensity exercises decreases the exercise-induced rise in plasma [K<sup>+</sup>] and enhances the concentration of Na<sup>+</sup>, K<sup>+</sup> pumps in trained human muscle by ~15%.<sup>[20-23]</sup> Thus McKenna MJ in his study pointed out that the physical training enhances, whilst inactivity hampers K<sup>+</sup> regulation while exercising.<sup>[20,22]</sup> Others workers have also opined that Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup>-transporter, K<sup>+</sup>-Cl<sup>-</sup> cotransporter, ATP-sensitive K<sup>+</sup> channel, and calcium-activated K<sup>+</sup> channel which are present in the skeletal muscle, and might contribute to extracellular potassium homeostasis in exercise.<sup>[21-23]</sup>

### **Exercise and Anti-Oxidant and Free Radical Levels**

Sen CK and Packer L while exploring thiol homeostasis opined that the physical exercise may induce oxidative stress and lead to blood glutathione oxidation.<sup>[25-27]</sup> The physical training programs and exercise protocols have specific effects on glutathione metabolism and it depends on the exercise protocol, duration and intensity of exercises. Glutathione metabolism in tissues sensitively responds to strenuous exercise. Similarly Gene expression of muscle mitochondrial (Mn) superoxide dismutase is enhanced after an acute bout of exercise and this is accompanied by increased level of NF-kappaB and AP-1 binding. The repeated bouts of exercise increases de novo protein synthesis of an antioxidant enzyme.<sup>[27-30]</sup>

Contraction-induced production of reactive oxygen species (ROS) produces oxidative stress to skeletal muscle. As an adaptive combat response the muscle antioxidant defence systems gets up regulated after heavy exercise. The nuclear factor (NF) kappaB and mitogen-activated protein kinases (MAPKs) are oxidative stress-sensitive signal transduction pathways in mammals. ROS may serve as messenger molecules and the adaptive responses act via redox-sensitive signalling pathways to maintain

cellular oxidant-antioxidant homeostasis in exercise.<sup>[31-35]</sup>

**Writing an Exercise Prescription**

This needs complete understanding of basic principles of energy expenditure and exercise pattern: Sequential understanding of principle are briefed point wise below.

1. Type of Exercises in practice in health and disease<sup>[30-32]</sup>
  - A. Aerobic exercise: These are endurance type exercise, rhythmic and sustained for sometimes. Example: Walking, jogging, running, cycling, swimming, etc.
  - B. Strength (Resistance) exercise: It includes weight training with free weight, machine, elastic rope, calisthenics, etc.
  - C. Flexibility exercise: These are stretching exercise.
2. Physical Activity Intensity in Metabolic Equivalent (MET)<sup>[30,34-37]</sup>
  - Metabolic Equivalent (MET) is the amount of energy expended during exercise relative to the energy expenditure during rest. (Note: Energy expenditure during rest = 1 MET; in Light Physical Activity: Less than 3 MET, Moderate Physical Activity: 3 – 6 MET and Vigorous Physical Activity: Above 6 MET)
3. Activities of Light, Moderate and Vigorous Physical Activities<sup>[35-38]</sup> are
  - A. Light Physical Activity: Leisure walking, cooking, daily household activities etc.
  - B. Moderate Physical Activities are: Brisk walking, Recreational swimming, Volleyball, Slow aerobics, Moderate cycling Gardening, Tennis-double, Badminton etc. Note that moderate Physical Activity Performed Regularly for 120 – 200 min. per week reduces the incidence of Diabetes.
  - C. Vigorous Physical Activities are: Jogging, Running, Tennis-single, Basketball, Rope skipping, Squash, Fast aerobics, Fast cycling, Stepping, Soccer, etc.
4. Energy Cost of Physical Activity
 

**Examples:** Slow Walking -2.5 MET, Fast walking 4MET, Swimming 6 MET, Tennis 8, Rope Skipping 8 MET, Basket Ball 8 MET etc.

**Calculating Energy Expenditure**

  - Energy Expenditure during Brisk Walking= 0.07 k. calorie per kg of body weight / min. An Example: a person weighing 80 kg would expend: 0.07 X 80 = 5.6 k. calorie per min.
  - If he has to expend 1200 k. calories per week.
    - ❖ How much time he should walk per week?
    - ❖  $1200 / 5.6 = 214.4$  min.
    - ❖ Duration of walking / day = 43 min / 5 days per week, or
    - ❖ Duration of walking / day = 54 min / 4 days per week.

**Exercise Prescription**

Exercise Regime in Overweight Individuals and Obesity without and clinical signs and symptoms includes.<sup>[37,39,40]</sup>

The moderate Intensity Physical Activity in which the energy expenditure = 3 - 6 MET and is practice for  $\geq 30$  min/day,  $\geq 5$  days/week to expend (Approximately 150 min. per week)and energy expended is  $\geq 1000$  k. calories/week with muscle strengthening activity for 2 or more days/week (notably 8-10 exercises/ 8-12 reps).

**Patients of Diabetes**<sup>[2,36,39]</sup>

- A. Aerobic activity for 30 min. increased up to 60 min. every day or most days/week.
- B. The heart rate during activity should be gradually increased to reach 60 – 70% of HR max.
- C. Moderate intensity weight training program should be done regularly to maintain muscle strength (8-12 repetitions 2 times /week).
- D. In patient is on insulin, he / she should avoid exercise if glucose levels below 100 mg/dl or above 250 mg/dl.
- E. Note that patients with patients above 35 years of age with long duration of history of diabetes coronary heart disease risk factors, micro vascular disease (retinopathy, nephropathy) and peripheral vascular disease should consult Physician before starting exercise protocol sessions.

**Exercise prescription in patients of hypothyroidism**<sup>[40-43]</sup>

The reduced levels of thyroid hormone affect body’s ability to burn calories at a normal rate. Hence shedding excess weight becomes difficult in hypothyroid patients. Patients who are overtly hypothyroid and on treatment with drugs the edema subsides over period of time but duration is unknown while severely hypothyroid and underweight patients may tend to gravitate towards the mean. Hypothyroidism causes fat build-up, which leads to insulin resistance in the body, leading up to weight gain and susceptibility to diabetes. Treatment for thyroid disorder is what benefits the patient in controlling weight while exercises act as placebos. Exercise and diet go hand in hand to manage weight for a hypothyroid patient. Exercise helps by sweating thus losing excess water and preventing fluid retention in the body. Exercise promotes better sleep patterns and relieves anxiety too.

**CONCLUSION**

Exercise is the best “medicine” for physical fitness and in achieving perfect sense of physical, social and mental wellbeing. All patients’ physical activity should be assessed. . Patients with cardiorespiratory diseases, metabolic syndromes, arrhythmia, acute pulmonary embolism, acute myocarditis, sickle cell

anemia, thalassemia etc. should consult Physician prior to following any exercise protocol. The moderate Intensity Physical Activity in which the energy expenditure = 3 - 6 MET and is practice for  $\geq 30$  min/day,  $\geq 5$  days/week to expend (Approximately 150 min. per week) and energy expended is  $\geq 1000$  k. calories/week with muscle strengthening and flexibility exercise should be followed for optimum health.

## REFERENCES

- Gibala MJ, Little JP, Van Essen M, Wilkin GP, burgomaster KA, Safdar A, Raha S, Tarnopolsky MA. Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *J Physiol.* 2006; 575: 901-911.
- Kowal P, Chatterji S, Naidoo N, Biritwum R, Fan W, et al. Data resource profile: the World Health Organization Study on global Ageing and adult health (SAGE). *Int J Epidemiol.* 2012; 41: 1639-1649.
- Maimoon L, Sultan C. Effect of physical activity on calcium homeostasis and calcitropic hormones: a review. *Calcific Tissue Int.* 2009 Oct; 85 (4):277-86.
- Henderson SA, Graham HK, Mollan RA, Riddoch C, Sheridan B, Johnston H. Calcium homeostasis and exercise. *Int Orthop.* 1989; 13 (1):69-73.
- Ljunghall S, Joborn H, Roxin LE, Skarfors ET, Wide LE, Lithell HO. Increase in serum parathyroid hormone levels after prolonged physical exercise. *Med Sci Sports Exerc.* 1988; 20:122-125.
- Rudberg A, Magnusson P, Larsson L, Joborn H. Serum isoforms of bone alkaline phosphatase increase during physical exercise in women. *Calcif Tissue Int.* 2000; 66:342-347.
- Guillemant J, Accarie C, Peres G, Guillemant S. Acute effects of an oral calcium load on markers of bone metabolism during endurance cycling exercise in male athletes. *Calcif Tissue Int.* 2004; 74:407-414.
- Tosun A, Bolukbasi N, Cingi E, Beyazova M, Unlu M. Acute effects of a single session of aerobic exercise with or without weight-lifting on bone turnover in healthy young women. *Mod Rheumatol.* 2006; 16:300-304.
- Kelly DE, Goodpaster BH. Effects of exercise on glucose homeostasis in Type 2 diabetes mellitus. *Med Sci Sports Exerc.* 2001 Jun; 33(6 Suppl):S495-501; discussion S528-9.
- Kelly D and Goodpaster BH explored in their study, the dose-response effect of exercise on improved glucose control in Type 2 DM and found that increasing levels of physical activity contribute to better diabetes prevention.
- Julie A Morgan, Frances Corrigan, and Bernhard T Baune. Effects of physical exercise on central nervous system functions: a review of brain region specific adaptations. *J Mol Psychiatry.* 2015; 3: 3.
- Greenwood BN, Foley TE, Day HEW, Campisi J, Hammack SH, Campeau S, et al. Freewheel Running Prevents Learned Helplessness/Behavioral Depression: Role of Dorsal Raphe Serotonergic Neurons. *J Neurosci.* 2003; 23(7):2889-98.
- Greenwood BN, Foley TE, Day HE, Burhans D, Brooks L, Campeau S, et al. Wheel running alters serotonin (5-HT) transporter, 5-HT1A, 5-HT1B, and alpha 1b-adrenergic receptor mRNA in the rat raphe nuclei. *Biol Psychiatry.* 2005; 57(5):559-68.
- Greenwood BN, Foley TE, Burhans D, Maier SF, Fleshner M. The consequences of uncontrollable stress are sensitive to duration of prior wheel running. *Brain Res.* 2005; 1033 (2):164-78.
- Greenwood BN, Strong PV, Dorey AA, Fleshner M. Therapeutic effects of exercise: wheel running reverses stress-induced interference with shuttle box escape. *Behav Neurosci.* 2007; 121(5):992-1000.
- Mischel NA, Llewellyn-Smith IJ, Mueller PJ. Physical (in) activity-dependent structural plasticity in bulbospinal catecholaminergic neurons of rat rostral ventrolateral medulla. *J Comp Neurol.* 2014; 522(3):499-513.
- Samman S (2012) Essentials of Human Nutrition. Zinc. In: Mann J, Truswell S (eds) (4th edn), Oxford University Press, UK 171-175.
- Overgaard K, Lindström T, Ingemann-Hansen T, Clausen T (2002) Membrane leakage and increased content of Na<sup>+</sup>-K<sup>+</sup> pumps and Ca<sup>2+</sup> in human muscle after a 100-km run. *J Appl Physiol* (1985) 92: 1891-1898.
- DeRuisseau KC, Chevront SN, Haymes EM, Sharp RG (2002) Sweat iron and zinc losses during prolonged exercise. *Int J Sport NutrExercMetab* 12: 428-437.
- McKenna MJ. *J Mol Cell Cardiol.* 1995 Apr; 27(4):941-9.
- Sejersted OM, Sjøgaard G. Dynamics and consequences of potassium shifts in skeletal muscle and heart during exercise. *Physiol Rev.* 2000; 80: 1411-1481.
- Sjøgaard G, Adams RP, Saltin B. Water and ion shifts in skeletal muscle of humans with intense dynamic knee extension. *Am J Physiol.* 1985; 248: R190-R196.
- Youn JH, McDonough AA. Recent advances in understanding integrative control of potassium homeostasis. *Annu Rev Physiol.* 2009; 71: 381-401.
- Sen CK, Packer L. Thiol homeostasis and supplements in physical exercise. *Am J Clin Nut.* 2000 Aug; 72(2 Suppl):653S-69S.
- Jill. Exercise-induced modulation of antioxidant defence. *Ann N Y Acad Sci.* 2002 Apr; 959:82-92.
- Constantinescu A, Han D, Packer L. Vitamin E recycling in human erythrocyte membranes. *J Biol Chem* 1993; 268:10906-13.
- Wardman P, von Sonntag C. Kinetic factors that control the fate of thiyl radicals in cells. *Methods Enzymol* 1995; 251:31-45.
- Kuzuya M, Naito M, Funaki C, Hayashi T, Asai K, Kuzuya F. Protective role of intracellular glutathione against oxidized low density lipoprotein in cultured endothelial cells. *Biochem Biophys Res Commun* 1989; 163:466-72.
- Barclay JK, Hansel M. Free radicals may contribute to oxidative skeletal muscle fatigue. *Can J Physiol Pharmacol* 1991; 69:279-84.
- Lew H, Quintanilha A. Effects of endurance training and exercise on tissue antioxidative capacity and acetaminophen detoxification. *Eur J Drug Metab Pharmacokinet* 1991; 16:59-68.
- Ji LL, Stratman FW, Lardy HA. Enzymatic down regulation with exercise in rat skeletal muscle. *Arch Biochem Biophys* 1988; 263:137-49.
- Kanter MM, Hamlin RL, Unverferth DV, Davis HW, Merola AJ. Effect of exercise training on antioxidant enzymes and cardiotoxicity of doxorubicin. *J Appl Physiol* 1985; 59:1298-303.
- Tiidus PM, Houston ME. Antioxidant and oxidative enzyme adaptations to vitamin E deprivation and training. *Med Sci Sports Exerc* 1994; 26:354-9.
- Tessier F, Margaritis I, Richard MJ, Moynot C, Marconnet P. Selenium and training effects on the glutathione system and aerobic performance. *Med Sci Sports Exerc* 1995; 27:390-6.
- Sen CK, Marin E, Kretzschmar M, Hanninen O. Skeletal muscle and liver glutathione homeostasis in response to training, exercise, and immobilization. *J Appl Physiol* 1992; 73:1265-72.
- Slentz CA, Tanner CJ, Bateman LA, Durham MT, Huffman KM, HOumard JA & Kraus WE. Effect of exercise training intensity on pancreatic  $\beta$  cell function. *Diabetes care.* 2009; 36:1807-1811.
- Nybo L, Sunstrup E, Jakobson MD, MOhr M, Hornstrup T, Simonsen L, Bulow J, Randers MB, Neilson JJ, Aagaard P

- &Krustrup P. High intensity training versus traditional exercise interventions for promoting health. *Med Sci Sports Exerc.* 2010;42: 1951-1958
38. Pinilla FG and Hillman C. The influence of exercise on cognitive abilities. *Compr Physiol.*2013;3(1): 403-428
  39. Burgomaster KA, Heigenhauser GJ &Gibala MJ. Effect of short term sprint interval training on human skeletal muscle carbohydrate metabolism during exercise and time trial performance. *J Appl Physiol.* 2006; 100: 2041-2047.
  40. Kehoe R, Wu SY, Leske MC, Chylack LT. Comparing self-reported and physician reported medical history. *American Journal of Epidemiology.* 1994; 139: 813–818.
  41. Ciloglu F, Peker I, Pehlivan A, Karacabey K, Ilhan N, Saygin O, Ozmerdivenli R. Exercise intensity and its effects on thyroid hormones. *Neuro Endocrinol Lett.* 2005 Dec;26(6):830-4.
  42. McKinnon RA, Orleans CT, Kumanyika SK, Haire-Joshu D, Krebs-Smith SM, et al. Considerations for an obesity policy research agenda. *American Journal of Preventive Medicine.* 2009; 36: 351–371. 39.
  43. Swinburn BA, Sachs G, Hall KD, McPherson K, Finegood DT, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet* 2011; 378: 804–814. 40.

**How to cite this article:** John N, John J. Exercise prescriptions in Obesity, Diabetes and Hypothyroidism: A Review. *Ann. Int. Med. Den. Res.* 2018; 4(3):PHXX-PHXX.

**Source of Support:** Nil, **Conflict of Interest:** None declared