

Dengue in North India: Teaching Hospital Based Study.

S. S. Patel¹, Bipin Shukla², Bhavana³, Dilip Ku. Singh Kushawaha⁴, Ravi Kumar Sharma⁴

¹Assistant Professor, Department of Microbiology, Hind Institute of Medical Sciences, Safedabad, Barabanki, U.P., India.

²Junior Resident, Department of Microbiology, Hind Institute of Medical Sciences, Safedabad, Barabanki, U.P., India.

³Senior Resident, Department of Microbiology, Hind Institute of Medical Sciences, Safedabad, Barabanki, U.P., India.

⁴Tutor, Department of Microbiology, Hind Institute of Medical Sciences, Safedabad, Barabanki, U.P., India.

ABSTRACT

Background: Dengue is a common arboviral mosquito-transmitted disease. Dengue fever is characterized by high fever of abrupt onset, sometimes with biphasic fever, severe headache, nausea, vomiting, ocular pain, cutaneous rash, muscles & joint pain. **Methods:** The patients were selected from attending in the Hind Institute of Medical Sciences, Safedabad, Barabanki, U.P., India. We included patients suffering from dengue fever & found positive NS1, IgM, IgG from August-2015 to October-2015. **Results:** Test analyzed by Rapid card test & automated analyzer. 12 patients were NS1, IgM, IgG positive & changes their haematological & biochemical values. **Conclusion:** We should control the arboviral transmitted mosquitoes (*Aedes aegypti*) which helps to prevent from dengue infection

Keywords: Dengue, Biphasic fever, Arbovirus & Mosquito.

INTRODUCTION

Dengue or “break bone fever” is a common arboviral mosquito-transmitted disease, causes of febrile illness in the tropical & subtropical region. Around 2.5 billion people are equally significant in tropical & subtropical region.^[1,2]

The name derived from the Swahili word for “the walk of a Dandie” in Spanish. The first case of the dengue fever was recorded during the Jin Dynasty (265-420 AD) I China. After the naming & identification of the disease in 1779 by Benjamin, first epidemic recognized simultaneously in Asia, Africa & North America in the 1780.^[3]

Dengue virus is a member of the Flavivirus family. There are four serotypes (antigenically) which are designated as DENV-1, DENV-2, DENV-3 & DENV-4.^[4] Arbovirus transmitted by *Aedes aegypti* & *Aedes albopictus* mosquitoes. Between 2006 & 2012 the National Vector Borne Disease Control Program reported an annual average 20,474 dengue cases & 132 deaths from dengue infection.^[5] In which include dengue fever, dengue haemorrhagic fever (DHF) & dengue shock syndrome (DSS).^[6-10]

Name & Address of Corresponding Author

Dilip Ku. Singh Kushawaha,
Tutor, Department of Microbiology,
Hind Institute of Medical Sciences, Safedabad, Barabanki,
U.P., India.
E-mail: dilipsinghnp@gmail.com

In India, an initial epidemic of dengue fever was reported in 1963-1964 on the Eastern region than reaching Delhi in 1967 & Kanpur in 1968. Simultaneously involved the southern part & gradually the whole country were involved with widespread epidemics of all the four serotypes of dengue virus co-circulating in India.^[11-21] In 1968, the epidemic in Kanpur due to DV-4 & 1969 during both DV-4 & DV-2. In 1966, Vellore DV-3

were isolated. In outbreaks of dengue occurred in Rajasthan by DV-1 & dv-3. DV-2 was the predominant serotype circulating in Northern India, Lucknow, Delhi & Gwalior while in 1997 DV-1 was isolated during the epidemic in Delhi.^[22-27]

Traditionally dengue fever characterized by high fever of abrupt onset, sometimes with biphasic fever, severe headache, nausea, vomiting, ocular pain, cutaneous rash, muscle & joint pain. In dengue haemorrhagic fever haemorrhagic signs, hemo-concentration, vascular leakage such as pleural effusion, ascites & hypoalbuminaemia which can progress to shock & death. Sign & symptoms of DHF with tachycardia or low pulse pressure (< 20 mm Hg) or hypotension (systolic blood pressure < 90 mm Hg), cold clammy skin & restlessness show in dengue shock syndrome.^[28]

According to WHO (2009) dengue categorized in dengue with or without warning signs or severe dengue. Dengue: fever & to these nausea, skin rash, vomiting, body ache, leucopenia, or any warning sign include abdominal pain, tenderness, persistent vomiting, fluid accumulation like effusions & ascites, bleeding, liver enlargement or rise in hematocrit with the rapid decrease platelet count. In severe dengue: plasma leakage, bleeding & organ impairment include hepatic transaminases elevated beyond 1000 IU/L & central nervous system manifestations like alteration in sensorium or cardiac or other organ involvement.^[29]

Liver enlargement, abdominal pain & anorexia are more common in dengue fever, but present in both dengue fever & dengue haemorrhagic fever. In the adult dengue patients the frequency of hepatomegaly & clinical jaundice ranges from 4%-52% range & 1.7%-17% simultaneously.^[30-33]

MATERIALS AND METHODS

The patients were selected from attending in the Hind Institute of Medical Sciences, Safedabad, Lucknow O.P.D./admitted to showing fever less than 2 weeks with or without body rashes, abdominal pain, vomiting & anorexia. All the patients were not suffering from dengue. We included 12 patients suffering from dengue fever & found positive NS1, IgM, IgG from August-2015 to October & remaining patients suffering from other diseases such as Malaria, Typhoid, So we excluded from this study. Patients were not graded according to WHO criteria. Blood samples collected in vacutainers for Hb, Haematocrit, Platelets, leukocyte count, AST, ALT & dengue viral specific antigen (NS1) & IgM, IgG antibody. Hb, Haematocrit, platelets count, leukocytes count done manually method/automated hematological-analyzer & AST, ALT values determined by the automated biochemistry - Analyzer. NS1, IgM, IgG

done by Rapid Card Test (manufactured by j. Mitra & Co. pvt, okhla ind. area, ph-1, New Delhi, India). The performance of the test was evaluated & compared with ELISA test (According to manufacturing laboratory manual).

RESULTS

We study on 110 patients which show clinical sign & symptoms as dengue but only 12 Patients shows NS1, 1Gg & IgM antibody positive. All the patients show body rash, vomiting, abdominal pain, back pain & splenomegaly. In hematological finding, the average of haemoglobin 12.0 gm/dl, haematocrit 38.29%, total leukocyte count 5.4 Lac. cells per cubic mm, thrombocyte count 1.11 lacks where mild decreases of platelet count. In biochemical finding, moderate decrease AST (191.33 IU/L) & ALT (232.25 IU/L) value in dengue patient's serum.

Table 1: showing haematological & biochemical parameters.

Investigation	Normal lab value	Range	Mean	Remarks
Haemoglobin (gm/dl)	M:13.0-18.0 F:12.0-15.0	9.0-14.6	12.00	<12.0 n=05 (41.66%)
Hamatocrit (%)	M:38.8-46.4 F:35.4-44.4	28.5-40.6	38.29	<39.0 n=05 (41.66%)
TLC /cubic mm	4000-11000	1800-16500	5.4	<4.0 n=04 (33.33%)
Platelet count (cells/cubic mm)	1.5-5.0 Lac.	30000-160000	1.11	<1.0 n=08 (66.66%)
ALT (IU/L)	0.8 - 4.5	31.0-1294	191.33	>45 n=12 (100%)
AST (IU/L)	10 - 40	59.0-1100	232.25	>40 n=12 (100%)

DISCUSSION

We have found various clinical presentations as fever, rash, headache, abdominal pain & few neurological manifestations. Rashes developed due to immunological mechanism- when the dengue virus inactivates with host cells, the release of cytokines & stimulation of the immunological mechanism by which vascular endothelial changes, infiltration of mono-nuclear cells & perivascular edema occurs. Retro-orbital pain & headache mostly from systemic inflammatory mediators & fever is non-specific immunological mechanism. Some neurological manifestations have also noted. Involvement of neurological manifestation in dengue due to neurotropism of dengue virus & immunological mechanisms. Dengue virus usually transmission during the rainy season when the temperature is conducive for

build-up of the vector population breeding in secondary habitats as well as for longer mosquito survival. This season is breeding season for mosquitoes & they needed protein for egg production. So small-size female mosquitoes are forced to take more blood meals to obtain protein. In our study, average platelet counts 1.11 Lac. cells per cubic mm & 66.66% cases show below than 100000 cells/cubic mm observed. The causes include dengue antigen binds to platelets & antibody mediated immunological destruction of platelets. A drop in platelet count may be occasionally observed in dengue haemorrhagic fever. Increase haematocrit value (in the same patient, age & sex) indicators the definitive evidence of increased vascular permeability & plasma leakage & fall in haematocrit was noticed during hospitalized patients were a febrile & rehydrated adequately. Decreases leukocyte count

in below 4000/cub mm in dengue infection may due to virus induced inhibition/destruction of myeloid progenitor cells, but in our study average leukocyte count 5.4 /cub mm observed.

Arise in serum transaminases (AST/ALT) observed in our study, which is more than normal ranges. The values of serum transaminase increased due to hepatic inflammation or liver involvement in dengue infection. Heparan sulphate plays a pivotal role for the intrusion of the dengue virus into the liver (Hep G2) cells. AST is more than ALT because the sources of AST are more such as heart, striated muscles, erythrocytes etc.

CONCLUSION

The dengue infection is major problems in India. The exact data are not available because many of the smaller outbreaks go unreported. So governments have to collect data properly, which helps to know the current status of dengue infection. The destruction of mosquito is necessary, which transmitted dengue virus. We should spray anti-mosquito pesticides for the destruction of the larval stage of *A. aegypti*, which helps to prevent from dengue infection.

REFERENCES

- Guzman MG, Halstead SB, Artsob H, Buchy P, Farrar J, Gubler GJ, Hunsperger E, Kroeger A, Margolis HS, Martinez E, et al. Dengue: a continuing global threat. *Nat Rev Microbiol.* 2010; 8:S7-16. (PubMed)
- Guzman MG, Kouri G. Dengue: an update. *Lancet Infect Dis.* 2002;2:33-42. (PubMed)
- Halstead SB. *Dengue (Tropical Medicine: Science & Practice)*. River Edge, N. J: Imperial College Press. 2008: 1-10.
- Khan E, Siddiqui J, Shakoor S, Mehraj V, Jamil B, Hasan R. Dengue outbreak in Karachi, Pakistan, 2006: experience at a tertiary care centre. *Trans R Soc Trop Med Hyg* 2007;101:1114-9.
- National Vector Borne Disease Control Programme. *Dengue Cases & Deaths in the Country since 2007*. Ministry of Health & Family Welfare, Directorate General of Health Services. Accessed May 30,2013.
- Beatty ME, Beutels P, Meltzer MI, Shepard DS, Hombach J, Hutubessy R, Dessis D, Coudeville L, Dervaux B, Wichmann O, Margolis HS, Kuritsky JN, 2011. Health economics of dengue: a systemic literature review & expert panel, s assessment. *Am J Trop Med Hyg.* 2013;84:473-488.
- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM, Brownstein JS, Scott TW, Farrar JJ, Hay SI. The global distribution & burden of dengue. *Nature.* 2013;496:504-507.
- Gubler DJ. Epidemic dengue/dengue hemorrhagic fever as a public health, social & economic problem in the 21st century. *Trends Microbiol.* 2002;10:100-103.
- Murray NE, Quam MB, Wilder-Smith A. Epidemiology of dengue: past, present & future prospectus. *Clin Epidemiol.* 2013;5:299-309.
- World Health Organization 2009. *Dengue: Guidelines for Diagnosis, Treatment, Prevention & control*. Geneva: World Health Organization.
- E. Myers RM, Reuben R, Rodrigues FM. Studies on dengue in vellor. South India. *Am J Trop Med Hyg.* 1966;15:580-7.
- Karamchandani PV. Dengue groups of fever in India. *Lancet* 1946;1:92.
- Ramakrishnan SP, Gelfand HM, Bose PN, Mukharjee RN. The epidemic of acute hemorrhagic fever, Calcutta, 163: epidemiological inquiry. *Indian J Med Res.* 1964;52:633-91.
- Chaudhuri RN, Saha TK, Chaudhuri AD. Dengue-like fever in Calcutta: further preliminary observations. *Bull Calcutta Sch Trop Med.* 1965;13:2-3.
- Krishnamurthy K, Kastury TE, Chittipantula G. Clinical & pathological studies o an outbreak of dengue-like illness in Visakhapatnam. *Indian J Med Res.* 1965;53:800-12.
- Paul SD, Dandawate CN, Banerjee K, Krishnamurthy K. Virological & serological studies on an outbreak of dengue-like illness in Visakhapatnam, Andrapradesh. *India J Med Res.* 1965;53:777-89.
- Balaya S, D lima LV, Pavri KM. Investigation on an outbreak of dengue in Delhi in 1967. *India J Med Res.* 1969;57:767-74.
- Chaturvedi UC, Mathur A, Kapoor AK, Mehrotra NK, Mehrotra RML. Virological study of an epidemiological of febrile illness with haemorrhagic manifestations at Knpur, India, during 1968. *Bull World Health Organ.* 1970;43:289-93.
- Chaturvedi UC, Kapoor AK, Mathur a, Chandra D, Khan AM, Mehrotra RML. A clinical epidemiological study of an epidemic of febrile illness with haemorrhagic manifestations which occurred at Kanpur, India in 1968. *Bull World Health Organ.* 1970;281-7.
- Myers RM, Carey DE, Banerjee K, Reuben R, Ramamurti DV. Recovery of dengue type 3 virus from human serum & *Aedes aegypti* in South India. *India J Med Res.* 1968;56: 781-7.
- Ghosh BN. Study on the epidemic of dengue-like fever in Pondicherry (1964-65 & 1965-66). *J India Med Assoc* 1968;51:261-4.
- Chaturvedi UC, Mathur A, Kapoor AK, Tandon HO, Mehrotra RML. Clinicovirological study of the recurrence of the dengue epidemic with haemorrhagic manifestation at Kanpur, during 1969. *Indian J Med Res.* 1972;60:329-33.
- Ghosh SN, pavri KR, Sheikh BH, D Lima LV, Mahadev PV, et al. Investigations on the outbreak of dengue fever in Ajmer City, Rajasthan State in 1969. Part 1. Epidemiological, Clinical & virological study of the epidemic. *Indian J Med Res.* 1974;62:511-22.
- Dar L, Broor S, Sengupta S, Xess I, Seth P. The first major outbreak of dengue hemorrhagic fever in Delhi, India. *Emerg Infect Dis.* 1999;5:589-90.
- Agrawal R, Kapoor S, Nagar R, Mishra A, Tandon R, et al. A clinical study of the patients with dengue hemorrhagic fever during the epidemic of 1996 at Lucknow, India. *Southeast Asian J Med Public Health.* 1999;30:735-40.
- Parida MM, Dash PK, Upadhaya C, Sexena P, Jana AM. Serological & virological investigation of an outbreak of dengue fever in Gwalior, India. *India J Med Res* 2002;116:248-54.
- Kurukumbi M, Wali JP, Broor S, Agrawal P, Seth P, Handa R, et al. Seroepidemiological & active surveillance of dengue fever/dengue haemorrhagic fever in Delhi. *Indian J Med Sci.* 2001;55:149-56.
- World Health Organization. *Dengue hemorrhagic fever: diagnosis, prevention & control*. 2nd ed. Geneva: WHO; 1997.
- WHO. *Dengue: guidelines for diagnosis, treatment, prevention & control*. Geneva, 2009.
- Prakash O, Almas A, Jafari SM, Akhtar J, Alishah H. Severity of acute hepatitis & its outcome in patients with dengue fever in a tertiary care hospital Karachi, Pakistan, (South Asia) *BMC Gastroenterol.* 2010;10:43. (PubMed)

31. Karoli R, Fatima J, Siddiqi Z, Kazmi KI, Sultania AR. Clinical profile of dengue infection at a teaching hospital in North India. *J Infect Dev Ctries.* 2012;6:551-554.
32. Saha AK, Maitra S, Hazra SC. Spectrum of hepatic dysfunction in 2012 dengue epidemic I Kolkata, West Bengal. *India J Gastroenterol.* 2013;32:400-403.
33. Trung DT, Thao le TT, Hien TT, Hung NT, Vinh NN, Hien PT, Chinh NT, Simmons C, Wills B. Liver involvement associated with dengue infection in adults in Vietnam. *Am J Trop Med Hyg.* 2010;83:774-780.

How to cite this article: Patel SS, Shukla B, Bhavana, Kushawaha DKS, Sharma RK. Dengue in North India: Teaching Hospital Based Study. *Ann. Int. Med. Den. Res.* 2016;2(2):158-61.

Source of Support: Nil, **Conflict of Interest:** None declared