

USG Guided Infraclavicular Perineural Infusion of Bupivacaine with Additives Adenosine and Magnesium Sulphate Compared for Upper Limb Surgeries.

Dharmalingam AL¹, Valli Sathyamoorthy², Kundhavi Devi Ranganathan,² Rajasekar R,³ Heber Anandan,⁴

¹Senior Assistant Professor, Department of Anaesthesiology of Government Kilpauk Medical College, Tamilnadu, India.

²Professor, Department of Anaesthesiology, Government Kilpauk Medical College, Affiliated to The Tamilnadu Dr. M.G.R. Medical University, Chennai, Tamilnadu, India.

³Senior Assistant Professor, Government Kilpauk Medical College, Kilpauk, Tamilnadu, India.

⁴Senior Clinical Scientist, Department of Clinical Research of Dr. Agarwal's Health care Limited, Tamilnadu, India.

Received: July 2016

Accepted: July 2016

Copyright: © the author(s), publisher. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Brachial plexus block has revolutionized the field of regional anesthesia for upper limb surgeries. Infraclavicular subcoracoid approach gives complete block without major complications. **Aim:** To compare the additives adenosine and magnesium sulphate with bupivacaine in infraclavicular infusion for upper limb surgeries and postoperative analgesia. **Methods:** 40 patients scheduled for elective unilateral upper limb surgeries involving distal arm/ elbow/ forearm/ hand divided into two groups A (n=20) and B (n=20) randomly. Group A - adenosine 6mgs with 28 ml 0.5% bupivacaine bolus followed by infusion of 0.25% bupivacaine 28ml with 6mgs of adenosine (2ml) at a rate of 5ml/hr. Group B - magnesium sulphate 75 mgs (in 2ml) with 28ml of 0.5% bupivacaine bolus followed by infusion of 0.25% bupivacaine 28ml with 75mgs of magnesium sulphate (in 2ml) at a rate of 5ml/hr in USG guided placement of infraclavicular catheter. **Results:** Statistical analysis showed Group A had a faster onset time of sensory and motor block and faster recovery when compared to group B. Group A needed more rescue analgesia than group B. Though Group B had a higher incidence of sedation and vomiting than group A, it was not significant. **Conclusion:** Though the addition of adenosine and magnesium sulphate to bupivacaine as additive gives prolonged analgesia, magnesium sulphate exceeds adenosine in achieving prolonged postoperative analgesia and better pain score and may be a better choice than adenosine.

Keywords: Bupivacaine, Adenosine, Magnesium Sulfate, Brachial plexus block.

INTRODUCTION

The infraclavicular brachial plexus block (ICB) is designed to deposit anesthetic high in the plexus, achieving anesthesia of the hand, forearm, elbow, and distal arm.^[1] Adenosine is a metabolic intermediate that is involved in nearly all aspects of cell function, including neurotransmission and signal transduction inside the body. Adenosine has an important role in the central and peripheral mediation of pain. Adenosine A1, A2A, A2B, and A3 receptors were detected in the spinal cord.

The A1 receptor plays an important role in spinal anti nociception where as the role of the A2A, A2B, and A3 receptors is still vague. A2A and A3 receptors mediate pain transmission peripherally, whereas the A1 receptor seems to have a central antinociceptive effect.^[2] Magnesium is the fourth most plentiful cation in the body and the second most plentiful intracellular cation after potassium. Anti-nociceptive effects of magnesium are due to regulation of calcium influx into the cell and antagonism of the N-methyl D-aspartate (NMDA) receptors. Many clinical investigations like Dogru et al^[3] and Choi et al^[4] have demonstrated that Mg administration in peripheral nerve plexus block has prolonged the analgesic effect and reduced postoperative analgesic consumption. Though magnesium has an analgesic property, it has not been studied well as an adjuvant to bupivacaine in infraclavicular brachial plexus block. The aim of this study to compare the additives adenosine and magnesium sulphate with bupivacaine

Name & Address of Corresponding Author

Dr. A. L. Dharmalingam
Department of Anaesthesiology
Kilpauk Medical College Hospital
Chennai – 600010
Tamilnadu, India.
alakshmanan33@yahoo.com

in infraclavicular infusion for upper limb surgeries and postoperative analgesia.

MATERIALS AND METHODS

A prospective comparative study was undertaken in 40 patients at Department of Anesthesiology, Kilpauk medical college hospital, Chennai. Institutional Ethics committee approval and informed consent were obtained. 20 patients in each groups belonging to 20 to 50 years of age with weight less than 70kgs from both gender and ASA status 1 and 2 are included. Exclusion criteria: Refusal, coagulopathy, neuropathy, injury and previous surgery of brachial plexus, allergy to drugs used, and technical difficulty are excluded. After securing an intravenous line and Boyles machine check, multichannel monitor connected to read pulse rate, NiBp, Spo2, emergency drugs ready, patient sedated with Inj. Midazolam 2mgs iv., Inj. ondansetron 4mgs iv, Inj. Ranitidine 50mgs iv. After scout scanning block performed under strict aseptic precautions. Observation: Sensory (using pinprick test, 3 point score) and motor (using modified Bromage scale, 3 point score) blockade assessed every two minutes after the block From time of completion of block to the complete loss of sensation (anaesthesia) of all the four nerves (radial, ulnar, median, musculocutaneous) is noted as sensory onset time and loss of finger movements is motor onset time. Recovery is regain of all sensations in all areas of four nerves as sensory and all movements of fingers as motor recovery. Visual analog score, sedation (Ramsay sedation score), nausea and vomiting score, need for rescue analgesics and postoperative complications were observed. Unsuccessful block requiring general anesthesia are excluded from the study. Pain in postoperative ward managed with iv paracetamol 1gm (if VAS>4) and Inj. Fentanyl (if VAS>6).

RESULTS

No difference observed in Age, Gender, Height, Weight and Duration of surgery. Pulse rate, NIBP, SpO2 didn't show any statistical difference between groups. Complete sensory block occurred at 8 minutes in group A and in group B the complete block occurred at 10 minutes. There was a statistically significant difference at 8 and 10 minutes between the two groups regarding onset time. Group A (P value=0.045) had a earlier onset than group B (P value=0.006). Group A recovered from sensory effect at 40 hours and group B at 44 hours completely. There was a statistically significant difference .Group A (P value 0.006) recovered earlier than group B (P value=0.002). Complete motor block occurred at 10 minutes in group A and group B took 14 minutes for complete motor block. There was a statistically significant difference between two groups at 10 minutes (P

value 0.006) and 14 minutes (P value=0.006). Group A recovered completely at 28 hours and group B at 32 hours. Group A (P value=0.002) and group B (P value=0.093) and it is statistically significant. There is a significant difference in pain (VAS) score from 4 minutes (P value <0.001) to 14minutes (P value 0.001) and from 28hrs (P value <0.0001) to 48 hrs (p value =0.011). Group A had pain relief at 8 minutes and B had at 10 minutes and group B had better pain score for prolonged time (44 hours) than group A (40 hours). Regarding sedation there was no statistically significant difference between two groups. Regarding nausea and vomiting score there was no statistically significant difference between two groups noted. Group B had more duration of analgesia (44 hours) than group A (40 hours) Group B needed lesser amount of rescue analgesics. No complications observed in both groups intraoperatively and postoperatively. Long term complications not observed due to patient poor compliance in follow up.

DISCUSSION

Infraclavicular subcoracoid brachial plexus block is deposition of local anaesthetic at the cord level where all the nerves supplying upper limb lie in a compact manner to get anaesthetized completely without much complications than other approach when performed under ultrasound guidance.^[1] The deposition of local anaesthetics alone cannot give prolonged duration of analgesia. Hence addition adjuvants are practiced. Here we added adenosine and magnesium sulphate as adjuvant and compared the effects of both drugs. Fukunga et al^[5] and Gan etal^[6] studied the pain reducing effect of adenosine in major surgeries by acting on adenosine receptors and found to be a good nonopioid analgesic in perioperative setting. Dogru et al, Choi et al^[3] used magnesium sulphate in their study and found magnesium sulphate gives postoperative analgesia by blocking NMDA receptors. The demographic profile is not statistically significant in our study as stated in Gunduz et al study and hemodynamic parameters also not showed any statistically significant difference between two groups.^[7] The sensory and motor onset time is early in group A than in group B. Recovery from sensory and motor effect is faster with group A than group B. Khaleed et al in their study concluded that adenosine has shorter onset time, lower mean VAS score over 48 hours.^[8] Ekmeckzi et al found the delayed onset and prolonged duration of analgesia when magnesium is given in femoral nerve block.^[9] Magnesium sulphate gives prolonged duration of analgesia and better pain scores than adenosine in our study. Dogru et al and Choi et al used magnesium in brachial plexus block and found prolonged analgesia and better pain scores.^[3,4] Kasturi et al^[10] in supraclavicular route, Lee et al in interscalene route and Gunduz et al^[7] in

axillary route in their study concluded the delayed onset and prolonged analgesic action and motor blockade of magnesium when used in brachial plexus block. The decreased duration of analgesic action of adenosine is due to rapid metabolism is stated in studies of Apan et al^[11] and Khaleed et al^[8]. They need continuous infusion for prolonged effect. Group B needed less rescue analgesics. Lee et al^[12] study also got the same result. Khaleed et al in his study claimed 48 hrs of good VAS score with infusion of 10ml/hr of adenosine 12mgs for two hours.^[8] We infused for six hours at a rate of 5ml/hr with 6mgs of adenosine. Sensory effect lasted for 44 hours. Sedation score; nausea and vomiting score did not show much difference between the two groups. Choi et al in their study has observed no significant difference regarding the same.^[13]

CONCLUSION

Continuous infusion of adenosine when compared to magnesium sulphate has shorter onset time for both sensory and motor blockade and recovers earlier from both effects. The magnesium sulphate takes a little longer time for its onset but gives prolonged duration of analgesia and better pain score postoperatively. Hence the addition of magnesium sulphate as an additive to bupivacaine in brachial plexus block may be a better choice when prolonged postoperative analgesia is required.

REFERENCES

1. Sandhu NS, Capan LM. Ultrasound-guided infraclavicular brachial plexus block. *Br J Anaesth.*2002;89:254-9.
2. Apan A, Ozcan S, Buyukkocak U, Anbarci O, Basar H. Perioperative intravenous adenosine infusion to extend postoperative analgesia in brachial plexus block. *Eur J Anaesthesiol.* 2003;20:916-9.
3. Dogru K, Yildirim D, Ulgey A, Aksu R, Bicer C, Boyaci A. Adding magnesium to levobupivacaine for axillary brachial plexus block in arteriovenous fistule surgery. *Bratisl Lek Listy.* 2012;113:607-9.
4. Choi IG, Choi YS, Kim YH, Min JH, Chae YK, Lee YK, et al. The Effects of Postoperative Brachial Plexus Block Using MgSO (4) on the Postoperative Pain after Upper Extremity Surgery. *Korean J Pain.* 2011;24:158-63
5. Fukunaga AF, Alexander GE, Stark CW. Characterization of the analgesic actions of adenosine: Comparison of adenosine and remifentanyl infusions in patients undergoing major surgical procedures. *Pain.*2003;101:129-38.
6. Gan TJ, Habib AS. Adenosine as a non-opioid analgesic in the perioperative setting. *Anesth Analg.*2007;105:487-94.
7. Gunduz A, Bilir A, Gulec S. Magnesium added to prilocaine prolongs the duration of axillary plexus block. *Reg Anesth Pain Med.* 2006;31:233-6.
8. Mohammed KM, Amnar AS. USG guided continuous infusion in infraclavicular block using bupivacaine alone and combined with adenosine for pain control in upper extremity surgeries. *Saudi J anaesth* 2011;5(2):132-37.
9. Ekmekci P, Bengisun ZK, Akan B, Kazbek BK, Ozkan KS, Suer AH. The effect of magnesium added to levobupivacaine for femoral nerve block on postoperative analgesia in patients undergoing ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2013;21:1119-24.
10. Mukherjee K, Das A, Roy S. Evaluation of magnesium as adjunct in ropivacaine induced supraclavicular brachial plexus block-a prospective, double blinded randomised study. *J Res Pharm Pract* 2014;3(4):123-129.
11. Apan A, Ozcan S, Buyukkocak U, Anbarci O, Basar H. Perioperative intravenous adenosine infusion to extend postoperative analgesia in brachial plexus block. *Eur J Anaesthesiol.* 2003;20:916-9.
12. Lee AR, Yi HW, Chung IS, Ko JS, Ahn HJ, Gwak MS, et al. Magnesium added to bupivacaine prolongs the duration of analgesia after interscalene nerve block. *Can J Anaesth.* 2012;59:21-7.
13. Choi IG, Choi YS, Kim YH, Min JH, Chae YK, Lee YK, et al. The Effects of Postoperative Brachial Plexus Block Using MgSO on the Postoperative Pain after Upper Extremity Surgery. *Korean J Pain.* 2011;24:158-63.

How to cite this article: Dharmalingam AL, Sathyamoorthy V, Ranganathan KD, Rajasekar R, Anandan H. USG Guided Infraclavicular Perineural Infusion of Bupivacaine with Additives Adenosine and Magnesium Sulphate Compared for Upper Limb Surgeries. *Ann. Int. Med. Den. Res.* 2016; 2(5):AN25-AN27.

Source of Support: Nil, **Conflict of Interest:** None declared